

SMALL TIME CHUNG TYPE LIL FOR LÉVY PROCESSES

FRANK AURZADA, LEIF DÖRING, AND MLADEN SAVOV

ABSTRACT. We prove Chung-type laws of the iterated logarithm for general Lévy processes at zero. In particular, we provide tools to translate small deviation estimates directly into laws of the iterated logarithm.

This reveals laws of the iterated logarithm for Lévy processes at small times in many concrete examples. In some cases, exotic norming functions are derived.

1. INTRODUCTION

A classical question in stochastic process theory is to understand the asymptotic behavior of a given stochastic process $X = (X_t)_{t \geq 0}$ on the level of paths. In the present work we consider general Lévy processes and find Chung type LIL (laws of the iterated logarithm) at zero, that is, given the Lévy process X , we aim at characterizing a norming function b satisfying

$$(1.1) \quad \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b(t)} = 1, \quad \text{where} \quad \|X\|_t := \sup_{0 \leq s \leq t} |X_s|.$$

The topic of large and small time fluctuations of Lévy processes has been studied extensively in the past (see for instance [10] for an overview and [3, 19]).

It is well-known that, via the Borel-Cantelli Lemma, Chung type LIL for a general stochastic process are connected to the so-called small deviation rate of the process, i.e.

$$(1.2) \quad -\log \mathbb{P}(\|X\|_t \leq \varepsilon), \quad \text{as } \varepsilon \rightarrow 0 \text{ and } t \rightarrow 0.$$

The main motivation for this paper originates from the recent work [2], where a framework for obtaining the small deviation rate (1.2) for general Lévy processes (but fixed t) is provided. The difficulty in passing over from the small deviation estimate to the respective LIL concerns circumventing the independence assumption of the Borel-Cantelli lemma.

In this paper we show how the asymptotics of (1.2) imply explicit LIL. We stress that it is not sufficient to have estimates for (1.2) for *fixed* t , which usually are referred to as small deviation estimates.

Small deviation problems are studied independently of LIL and have connections to other fields, such as the approximation of stochastic processes, coding problems, the path regularity of the process, limit laws in statistics, and entropy numbers of linear operators. We refer to the surveys [12, 13] for an overview of the field and to [14] for a regularly updated list of references, which also includes references to laws of the iterated logarithm of Chung type. The papers [25, 18, 7, 21, 22, 16, 15, 17, 23, 24] provide a good source for earlier results on small deviations of Lévy processes.

We now discuss LIL for special Lévy processes that have already appeared in the literature. The norming function $b(t) = \sqrt{\pi^2 t / (8 \log |\log t|)}$ for a standard Brownian motion can be derived from the large time LIL, proved by Chung [9], via time inversion.

For any Lévy process with non-trivial Brownian component the recent result of [8] shows that

Date: January 6, 2011.

2000 Mathematics Subject Classification. Primary 60G51; Secondary 60F15.

Key words and phrases. Law of the Iterated Logarithm; Lévy Process; Small Ball Problem; Small Deviations.

The first author was supported by the DFG Emmy Noether Programme.

The second author was supported by the EPSRC grant EP/E010989/1.

The third author acknowledges the support of New College, Oxford.

(1.1) holds with the norming function for a standard Brownian motion. If X is an α -stable Lévy process (1.1) holds with norming function $b(t) = (c_\alpha t / \log |\log t|)^{1/\alpha}$, which goes back to [25].

Of course it is natural to ask for the general structure of the norming function for arbitrary Lévy processes not having the special features of the examples mentioned so far. LIL for more general Lévy processes were obtained by Wee in [26] (see [27] for more examples). It was shown that if for some positive constant θ

$$(1.3) \quad \mathbb{P}(X_t > 0) \geq \theta \quad \text{and} \quad \mathbb{P}(X_t < 0) \geq \theta, \quad \text{for all } t \text{ sufficiently small,}$$

holds, then upper and lower bounds in the LIL hold in the following sense: for λ_1 sufficiently small and λ_2 sufficiently large,

$$1 \leq \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda_1}(t)} \quad \text{and} \quad \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda_2}(t)} \leq 1$$

for norming functions b_λ given by

$$b_\lambda(t) := f^{-1} \left(\frac{\log |\log t|}{\lambda t} \right),$$

where f is given by some explicit, but complicated expression depending on the Lévy triplet.

Although the results of Wee are quite general, there are some drawbacks which we aim to overcome in the present work.

First, the proofs given in [26] and [27] are difficult to follow. This might be due to the use of the old-fashioned notation for the Lévy measure. In particular, it does not become clear that actually the LIL follow from small deviation estimates of type (1.2) and which behavior of the process is actually responsible for the correct norming function.

Secondly, even in the symmetric case, if the norming function b_λ is not regularly varying at zero, the unspecified (and suboptimal) constants λ_1 and λ_2 do not only appear in a weaker limiting constant, but they influence the norming function essentially (see (3.2) below for an example of influence on the exponential level). In our approach, we keep track of the appearing constants in an optimal way. In particular, in the case of known strong small deviation order, our approach allows to transfer the constant in the strong small deviation order to limiting constant in the LIL. Thirdly, although condition (1.3) looks innocent at a first glance it turns out to be quite delicate. It is certainly fulfilled for symmetric Lévy processes. Unfortunately, only given the Lévy triplet it seems to be unknown how (1.3) can be checked. On the contrary, our conditions are explicit in terms of the Lévy triplet. We give a couple of examples showing that our conditions can be checked easily.

This paper is structured as follows. In Section 2, we give the main results that manage the transfer between small deviations and LIL. Several examples of LIL for concrete Lévy processes are collected in Section 3. The proofs are given in Section 4.

Let us finally fix some notation. In this paper we let X be a Lévy process with characteristic triplet (γ, σ^2, Π) , where $\gamma \in \mathbb{R}$, $\sigma^2 \geq 0$, and the Lévy measure Π has no atom at zero and satisfies

$$\int (1 \wedge x^2) \Pi(dx) < \infty.$$

For basic definitions and properties of Lévy processes we refer to [3, 19]. As we are interested only in the behavior for small times we may truncate large jumps. In particular, we restrict ourselves to Lévy processes involving only jumps of absolute value at most 1. Hence, the characteristic exponent, $\mathbb{E}e^{izX_t} =: e^{t\psi(z)}$, has the form

$$\psi(z) = i\gamma z - \frac{\sigma^2 z^2}{2} + \int_{-1}^1 (e^{izx} - 1 - izx) \Pi(dx), \quad z \in \mathbb{R}.$$

For later use we denote by Φ the Laplace exponent of a subordinator A , $\mathbb{E}e^{-uA_1} = e^{-\Phi(u)}$,

$$\Phi(u) = u\gamma_A + \int_0^\infty (1 - e^{-ux}) \Pi_A(dx).$$

Further, we use the standard notation $\bar{\Pi}(\varepsilon) := \Pi([- \varepsilon, \varepsilon]^c)$ for the two-sided tail of the Lévy measure.

In the following, we denote by $f \sim g$ the strong asymptotic equivalence, i.e. $\lim f/g = 1$, and by $f \approx g$ the weak asymptotic equivalence, i.e. $0 < \liminf f/g \leq \limsup f/g < \infty$.

2. MAIN RESULTS

Our first theorem manages the transfer from small deviation rates to LIL under minimal loss of constants.

Theorem 1. *Let X be a Lévy process and F be a function increasing to infinity at zero such that with some $0 < \lambda_1 \leq \lambda_2 < \infty$*

$$(2.1) \quad \lambda_1 F(\varepsilon)t \leq -\log \mathbb{P}(\|X\|_t < \varepsilon) \leq \lambda_2 F(\varepsilon)t, \quad \text{for all } \varepsilon < \varepsilon_0 \text{ and } t < t_0.$$

Further, define

$$b_\lambda(t) := F^{-1}\left(\frac{\log |\log t|}{\lambda t}\right)$$

for $\lambda > 0$ and assume that, as $n \rightarrow \infty$,

$$(2.2) \quad (n+1)^{-(n+1)^\beta} \left| \int_{|x| > b_{\lambda'_2}(n^{-n^\beta})} x \Pi(dx) - \gamma \right| = o(b_{\lambda'_2}(n^{-n^\beta})), \quad \text{for all } \beta > 1 \text{ and } \lambda'_2 > \lambda_2.$$

Then the LIL

$$1 \leq \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda'_1}(t)} \quad \text{and} \quad \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda'_2}(t)} \leq 1$$

hold almost surely for any $\lambda'_1 < \lambda_1$ and $\lambda'_2 > \lambda_2$.

Remark 1. *It is important to note the role of (2.2). It ensures that the process does not become too asymmetric when one continues to cut off more and more smaller jumps. Only in this case it is possible to expect an estimate of type (2.1) to follow from the framework given in [2]. Corollary 4 below and in particular (2.11) give a sufficient condition when this is the case.*

Remark 2. *It may be instructive to study (2.2) in the case of a particular example. Namely, one can easily check that (2.2) holds when we have a stable subordinator. Moreover by simple computation using the fact that $\Pi(dx) = cx^{-\alpha-1}dx$ and Theorem 11 in [3] we get that the right norming function in this case is*

$$b(t) = t \left| \int_{x>t} x \Pi(dx) \right|,$$

which is almost the left-hand side of (2.2). This shows that (2.2) is not of technical matter, but the quantities arising there can acutally govern the norming function in the LIL, in which case Theorem 1 cannot be applied.

Remark 3. *Let us compare (2.2) with the condition from Wee [26]. Note that condition (2.2) is analytic, i.e. in terms of the Lévy triplet. Contrary, Wee's condition (1.3) is easily formulated and has a particular probabilistic interpretation. But unfortunately there exists no necessary and sufficient condition in the terms of the Lévy triplet which allows for the verification of (1.3). To understand the difficulty behind obtaining such a condition it may be instructive to look at Theorems 4 and 5 in [1], which reformulate (1.3) in terms of other probabilistic quantities.*

It is crucial that there is almost no loss of constants in the transfer from the small deviations to the LIL as in cases when b_λ is not regularly varying, the constants λ'_1, λ'_2 may influence the rate function drastically (see (3.2) for an extreme example).

If instead b_λ only depends on λ via a multiplicative constant, our approach allows to strengthen the previous theorem to the optimal limiting constants. Such examples occur for instance if the small deviation rate function F is regularly varying.

Corollary 1. *In the setting of Theorem 1 assume additionally that F is regularly varying at zero with non-positive exponent. Then the following LIL hold almost surely:*

$$(2.3) \quad 1 \leq \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda_1}(t)} \quad \text{and} \quad \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda_2}(t)} \leq 1.$$

In particular, if there is $\lambda > 0$ such that (2.1) holds for all $\lambda_1 < \lambda$ and all $\lambda_2 > \lambda$ then

$$\liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_\lambda(t)} = 1 \quad \text{a.s.}$$

In the setting of a regularly varying rate function, say F is regularly varying at zero with exponent $-\alpha$, $\alpha > 0$, one can express (2.3) as

$$\liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_1(t)} \in [\lambda_1^{1/\alpha}, \lambda_2^{1/\alpha}], \quad \text{a.s.}$$

This shows that only the quality of the small deviation estimate (2.1) matters in order to obtain the limiting constant in the LIL. Recall that the Blumenthal zero-one law implies that the limit is almost surely equal to a deterministic constant, which in this case can be specified.

Theorem 1 reduces the question of the right norming function for the LIL to the question of small deviations which is known precisely for many examples. For general Lévy processes those have been obtained in [2] (their results were stated for $t = 1$ only but hold in general as we discuss in Proposition 1 below). In particular, for symmetric Lévy processes their main result states that the rate function is given by

$$(2.4) \quad F(\varepsilon) = \varepsilon^{-2}U(\varepsilon),$$

where $U(\varepsilon)$ is the variance of X with jumps larger than ε replaced by jumps of size ε :

$$(2.5) \quad U(\varepsilon) := \varepsilon^2 \bar{\Pi}(\varepsilon) + \sigma^2 + \int_{-\varepsilon}^{\varepsilon} x^2 \Pi(dx).$$

From these specific small deviations we can deduce the following corollary for symmetric processes.

Corollary 2. *Let X be a symmetric Lévy process, then there are $0 < \lambda_1 \leq \lambda_2 < \infty$ such that almost surely,*

$$1 \leq \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda_1}(t)} \quad \text{and} \quad \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda_2}(t)} \leq 1,$$

with

$$b_\lambda(t) := F^{-1} \left(\frac{\log |\log t|}{\lambda t} \right)$$

and F defined in (2.4). If additionally F is regularly varying at zero with exponent $-\alpha$, $\alpha > 0$, then the following general bounds hold:

$$\frac{1}{12} \frac{1}{2^\alpha} \leq \lambda_1 \leq \lambda_2 \leq 3^\alpha 10.$$

The loss of constants in the corollary is only due to the general formulation. For some examples we will see below that the small deviations are known in the strong asymptotic sense so that Theorem 1 gives the precise law.

For “strongly non-symmetric” Lévy processes we have to proceed differently, since here condition (2.2) does not hold. For this case, we provide another, different link between small deviation rates and LIL; we keep track of the constants in the norming function in an optimal way and only lose the limiting constant.

Theorem 2. *Let X be a Lévy process and F be a function increasing to infinity at zero such that for $0 < \lambda_1 \leq \lambda_2 < \infty$*

$$(2.6) \quad \lambda_1 F(\varepsilon)t \leq -\log \mathbb{P}(\|X\|_t < \varepsilon) \leq \lambda_2 F(\varepsilon)t, \quad \text{for all } \varepsilon < \varepsilon_0 \text{ and } t < t_0.$$

Furthermore, set

$$(2.7) \quad b_\lambda(t) := F^{-1} \left(\frac{\log |\log t|}{\lambda t} \right)$$

and suppose that there is a constant $C > 0$ such that

$$(2.8) \quad C b_\lambda(t) \leq b_\lambda(t/2), \quad 0 < t \leq t_0, \lambda \in (\lambda_1/2, 2\lambda_2).$$

Then the LIL

$$0 < \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda'_1}(t)} \quad \text{and} \quad \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda'_2}(t)} < \infty$$

hold almost surely for all $\lambda'_1 < \lambda_1$ and $\lambda_2 < \lambda'_2$.

Again, if the rate function F is regularly varying, then we can strengthen the result.

Corollary 3. *In the setting of Theorem 2 assume additionally that F is regularly varying at zero with negative exponent. Then the following LIL hold almost surely:*

$$\liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_1(t)} \in (0, \infty).$$

The theorems listed so far manage the transfer between small deviation order and LIL. Similarly to Corollary 2, we can combine them with the main results of [2]. This looks more technical in the present case. We give an explanation of the role of the different terms after stating the result.

Corollary 4. *Let X be a Lévy process with triplet (γ, σ^2, Π) . Assume that u_ε is the solution of the equation $\Lambda'_\varepsilon(u) = 0$, where Λ_ε is the following log Laplace transform:*

$$(2.9) \quad \Lambda_\varepsilon(u) = \frac{\sigma^2}{2} u^2 + \left(\gamma - \int_{[-1,1] \setminus [-\varepsilon, \varepsilon]} x \Pi(dx) \right) u + \int_{-\varepsilon}^{\varepsilon} (e^{ux} - 1 - ux) \Pi(dx).$$

Set

$$(2.10) \quad F(\varepsilon) := \varepsilon^{-2} U_\varepsilon(\varepsilon) - \Lambda_\varepsilon(u_\varepsilon), \quad U_\varepsilon(\varepsilon) := \varepsilon^2 \bar{\Pi}(\varepsilon) + \sigma^2 + \int_{-\varepsilon}^{\varepsilon} x^2 e^{-u_\varepsilon x} \Pi(dx),$$

and assume F is increasing to infinity as $\varepsilon \rightarrow 0$. Define b as in (2.7) and assume that b satisfies (2.8). If furthermore

$$(2.11) \quad \varepsilon |u_\varepsilon| = o(\log \log F(\varepsilon)), \quad \text{as } \varepsilon \rightarrow 0,$$

is satisfied then we have for some $\lambda_1, \lambda_2 > 0$

$$0 < \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda_1}(t)} \quad \text{and} \quad \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda_2}(t)} < \infty \quad \text{a.s.}$$

Let us explain the quantities occurring in Corollary 4 in more detail. The main observation is that the proof for the small deviation estimates in [2] (Theorem 1.5) can be used directly for any $t > 0$ to obtain the following proposition.

Proposition 1. *Let Λ_ε be as defined in (2.9) and assume that u_ε is the solution of $\Lambda'_\varepsilon(u_\varepsilon) = 0$. Then, with F as in (2.10), we have for all $t > 0$ and all $\varepsilon < 1$*

$$(2.12) \quad \frac{1}{12} t F(2\varepsilon) - \varepsilon |u_{2\varepsilon}| - 1 \leq -\log \mathbb{P}(\|X\|_t \leq \varepsilon) \leq 10t F\left(\frac{\varepsilon}{3}\right) + \varepsilon |u_{\varepsilon/3}| + 3.$$

The term $\bar{\Pi}(\varepsilon)$ in (2.12) (included in the F term) comes from the requirement that there should be no jumps larger than ε . After removing these jumps, the process may drift out of the interval $[-\varepsilon, \varepsilon]$, which is prevented by applying an Esscher transform to the process, whose ‘price’ is given by the term $-\Lambda_\varepsilon(u_\varepsilon)$. The quantity u_ε is the drift that has to be subtracted in order to make the process a martingale. Then the remaining process is treated as in the symmetric case, and the same term $\varepsilon^{-2} U_\varepsilon(\varepsilon)$ appears as in (2.4), but this time with respect to the Lévy measure transformed by the change of measure.

Note that (2.12) is almost the required estimate in (2.6), except for the term $\varepsilon |u_\varepsilon|$, which may spoil the estimate. It is exactly condition (2.11) that ensures that the term $\varepsilon |u_\varepsilon|$ can be neglected.

We stress that in some cases $\varepsilon|u_\varepsilon|$ does give an order that is larger than $tF(\varepsilon)$ so that the function b from (2.7) is not the right norming function. This effect can be observed in some examples below. In particular, this happens for processes of bounded variation with non-zero drift.

Proposition 2. *Let X be a Lévy process with bounded variation and non-vanishing effective drift, i.e. $\int_{[-1,1]} |x|\Pi(dx) < \infty$ and $c := \gamma - \int_{-1}^1 x \Pi(dx) \neq 0$. Then*

$$\lim_{t \rightarrow 0} \frac{\|X\|_t}{t} = |c| \quad a.s.$$

The proof of this proposition is based on classical arguments rather than any connection to small deviations.

3. EXPLICIT LIL FOR LÉVY PROCESSES

In this section we collect concrete Lévy processes for which we can transform small deviation results to an LIL. As we have seen, understanding the small deviation rates is crucial.

The first corollary gives us a useful variance domination principle for LIL that works for many examples.

Corollary 5. *Suppose X^1 and X^2 are independent symmetric Lévy processes, then $X^1 + X^2$ and X^2 fulfill precisely the same LIL if*

$$\lim_{\varepsilon \rightarrow 0} \frac{U_{X^1}(\varepsilon)}{U_{X^2}(\varepsilon)} = 0.$$

Proof. This follows directly from Corollary 2 noticing that $U_{X^1+X^2} = U_{X^1} + U_{X^2}$. \square

In the same spirit the following corollary (recovering (3.2) in [8]) displays the intuitive fact that a non-zero Brownian component dominates the jumps of a Lévy process.

Corollary 6. *If X is a Lévy process with $\sigma \neq 0$, then*

$$\liminf_{t \rightarrow 0} \frac{\|X\|_t}{\sqrt{t/\log|\log t|}} = \frac{\pi\sigma}{\sqrt{8}} \quad a.s.$$

Proof. Following precisely the proof of Corollary 2.6 of [2] one can show that the small deviation rates of Lévy processes with non-zero Brownian component is given by

$$-\log \mathbb{P}(\|X\|_t < \varepsilon) \sim \frac{\pi^2 \sigma^2}{8} \varepsilon^{-2} t, \quad \text{as } \varepsilon \rightarrow 0 \text{ and } t \rightarrow 0.$$

Hence, the norming function follows from Theorem 1. As the process is not necessarily symmetric, condition (2.2) has to be checked: Since $b(t) = \sqrt{t\pi^2/(8\log|\log t|)}$ and $\int_{|x|>\varepsilon} |x|\Pi(dx) = o(\varepsilon^{-1})$, it remains to be seen that

$$a_{n+1} \leq cb(a_n)^2 = a_n/\log| \log a_n|$$

for $a_n = n^{-n^\beta}$ and $\beta > 1$. This can be verified by simple computations. \square

Similarly to Lévy processes with non-zero Brownian component, symmetric processes of smaller small deviation order (e.g. stable processes of smaller index) are dominated by stable Lévy processes.

Corollary 7. *Let X be a symmetric α -stable Lévy process with $\alpha \in (0, 2]$ and let Y be symmetric with $U_Y(x) = o(x^{2-\alpha})$. Then there is a constant $0 < c_\alpha < \infty$ such that*

$$\liminf_{t \rightarrow 0} \frac{\|X+Y\|_t}{(t/\log|\log t|)^{1/\alpha}} = \liminf_{t \rightarrow 0} \frac{\|X\|_t}{(t/\log|\log t|)^{1/\alpha}} = c_\alpha^{1/\alpha} \quad a.s.$$

Proof. The small deviation rate is given by

$$-\log \mathbb{P}(\|X\|_t < \varepsilon) \sim c_\alpha \varepsilon^{-\alpha} t, \quad \text{as } \varepsilon \rightarrow 0 \text{ and } t \rightarrow 0,$$

for some constant $c_\alpha > 0$ (see e.g. page 220 in [3]). Hence, the LIL follow from Corollary 1 and Corollary 5. \square

Remark 4. The constant c_α in the LIL of stable Lévy processes is the unknown constant of the small deviations for respective α -stable Lévy processes (see [25] and Proposition 3 and Theorem 6 in Chapter VIII of [3]). The results of [2] entail the following concrete bounds:

$$\frac{2C}{2^\alpha} \left(\frac{1}{\alpha} + \frac{1}{12(2-\alpha)} \right) < c_\alpha < 3^\alpha \cdot 2C \left(\frac{1}{\alpha} + \frac{10}{2-\alpha} \right),$$

where C is the constant in the Lévy measure: $\Pi(dx) = C|x|^{-(1+\alpha)}dx$. This implies $c_\alpha \sim 2C/\alpha$, as $\alpha \rightarrow 0$. We remark that, contrary to the symmetric case, the constant c_α is known explicitly for completely asymmetric stable Lévy processes, see [4].

Let us study the case that Π behaves as a regularly varying function at zero and is symmetric. Then the following LIL are satisfied.

Corollary 8. Let X be a Lévy process with triplet $(0, 0, \Pi)$ with Π being symmetric and

$$\bar{\Pi}(\varepsilon) \approx \varepsilon^{-\alpha} |\log \varepsilon|^{-\gamma}, \quad \text{as } \varepsilon \rightarrow 0,$$

with $0 < \alpha < 2$ or $\alpha = 2, \gamma > 1$. Then

$$\liminf_{t \rightarrow 0} \frac{\|X\|_t}{b(t)} \in (0, \infty) \quad \text{a.s.}$$

with

$$b(t) = \begin{cases} \left(\frac{t |\log t|^{-\gamma}}{\log |\log t|} \right)^{1/\alpha} & : 0 < \alpha < 2, \\ \left(\frac{t |\log t|^{1-\gamma}}{\log |\log t|} \right)^{1/2} & : \alpha = 2, \gamma > 1. \end{cases}$$

Proof. The corollary follows from Theorem 1. The required small deviation estimate,

$$-\log \mathbb{P}(\|X\|_t < \varepsilon) \approx \begin{cases} \varepsilon^{-\alpha} |\log \varepsilon|^{-\gamma} t & : 0 < \alpha < 2, \\ \varepsilon^{-2} |\log \varepsilon|^{1-\gamma} t & : \alpha = 2, \gamma > 1, \end{cases}$$

as $\varepsilon \rightarrow 0$ and $t \rightarrow 0$, is obtained from Proposition 1 (cf. Example 2.2 in [2] for $t = 1$). Since we deal with a symmetric process, condition (2.11) is trivially satisfied due to $u_\varepsilon = 0$. \square

Having discussed the α -stable like cases, we now consider Lévy processes with polynomial tails near zero of *different* exponents. The technique used for this example can be extended to any case with essentially regularly varying Lévy measure at zero. Let X be a Lévy process with triplet $(\gamma, 0, \Pi)$, where Π is given by

$$(3.1) \quad \frac{\Pi(dx)}{dx} = \frac{C_1 \mathbf{1}_{(0,1]}(x)}{x^{1+\alpha_1}} + \frac{C_2 \mathbf{1}_{[-1,0)}(x)}{(-x)^{1+\alpha_2}},$$

with $2 > \alpha_1 \geq \alpha_2$ and $C_1, C_2 \geq 0$, $C_1 + C_2 \neq 0$. We now analyze the pathwise behavior at zero in the cases when $\alpha_1 > 1$, $\alpha_1 = 1$, and $0 < \alpha_1 < 1$, respectively. The second exponent α_2 can be even negative.

Corollary 9. Let X be a Lévy process with triplet $(\gamma, 0, \Pi)$ with Π as in (3.1). Then the following holds:

(1) If $\alpha_1 \geq \alpha_2$, $C_1 \neq 0$, and $\alpha_1 > 1$, then

$$\liminf_{t \rightarrow 0} \frac{\|X\|_t}{(t/\log |\log t|)^{1/\alpha_1}} \in (0, \infty) \quad \text{a.s.}$$

(2) If $\alpha_1 = \alpha_2 = 1$ and $C_1 = C_2$ then

$$\liminf_{t \rightarrow 0} \frac{\|X\|_t}{t/\log |\log t|} \in (0, \infty) \quad \text{a.s.}$$

(3) If $1 > \alpha_1 \geq \alpha_2$ and the effective drift does not vanish, then

$$\lim_{t \rightarrow 0} \frac{\|X\|_t}{t} = |c| \quad \text{a.s.}$$

Proof. Parts (1) and (2) follow from Theorem 1. The required small deviation estimates,

$$-\log \mathbb{P}(\|X\|_t < \varepsilon) \approx \varepsilon^{-\alpha_1 t}$$

for $\varepsilon \rightarrow 0$ and $t \rightarrow 0$, are obtained from Proposition 1 (cf. Corollary 2.7, 2.8, and 2.9 of [2] for $t = 1$; note that $u_\varepsilon \approx \varepsilon^{-1}$ in all cases). One can easily check condition (2.11).

In part (3), the process is of bounded variation, so that the claim is included in Proposition 2. \square

We now come to Lévy processes obtained from Brownian motion by subordination, i.e. $X_t = \sigma B_{A_t}$, where B is a Brownian motion independent of the subordinator A . In this case, the resulting Lévy process is symmetric and the small deviation asymptotics is governed by the truncated variance U from (2.5).

Corollary 10. *Let B be a Brownian motion independent of the subordinator A , where A has Laplace exponent Φ . For $\lambda > 0$ we set $b_\lambda(t) := F^{-1}\left(\frac{\log |\log t|}{\lambda t}\right)$ with*

$$F(\varepsilon) := \Phi(\sigma^2 \varepsilon^{-2}) + \gamma_A \sigma^2 \varepsilon^{-2}.$$

Then for some $\lambda_1, \lambda_2 > 0$

$$1 \leq \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda_1}(t)} \quad \text{and} \quad \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda_2}(t)} \leq 1 \quad a.s.$$

In particular, if $\gamma_A = 0$ and Φ is regularly varying with positive exponent, we have

$$\liminf_{t \rightarrow 0} \frac{\|X\|_t}{(\Phi^{-1}(\log |\log t|/t))^{-1/2}} \in (0, \infty) \quad a.s.$$

Proof. The corollary follows from Theorem 1 with the small deviation estimate from Proposition 1

$$-\log \mathbb{P}(\|X\|_t \leq \varepsilon) \approx (\Phi(\sigma^2 \varepsilon^{-2}) + \gamma_A \sigma^2 \varepsilon^{-2})t,$$

as $\varepsilon \rightarrow 0$ and $t \rightarrow 0$ (cf. Example 2.13 of [2] for $t = 1$ and note the misprint there). Condition (2.2) is trivially fulfilled as the process is symmetric. \square

For a more specific example, in particular exhibiting exotic small time behavior, we choose the subordinator A to be a Gamma process. Then one defines the so called Variance-Gamma process as

$$X_t = \sigma B_{A_t} + \mu A_t,$$

for some constants $\sigma \neq 0$ and $\mu \in \mathbb{R}$.

Corollary 11. *Let X be a Variance-Gamma process, then for $\mu = 0$ there are some constants $0 < \lambda_1 \leq \lambda_2 < \infty$ such that*

$$(3.2) \quad 1 \leq \liminf_{t \rightarrow 0} \frac{\|X\|_t}{e^{-\lambda_1 \log |\log t|/t}} \quad \text{and} \quad \liminf_{t \rightarrow 0} \frac{\|X\|_t}{e^{-\lambda_2 \log |\log t|/t}} \leq 1 \quad a.s.,$$

whereas for $\mu \neq 0$

$$\liminf_{t \rightarrow 0} \frac{\|X\|_t}{t} = |\mu| \mathbb{E}A_1 \quad a.s.$$

Proof. The second part is included in Proposition 2, since the process is of bounded variation with non-zero effective drift. In the first part, the effective drift is zero, and the claim follows from Theorem 1. The small deviation estimate,

$$-\log \mathbb{P}(\|X\|_t \leq \varepsilon) \approx t |\log \varepsilon|, \quad \text{as } \varepsilon \rightarrow 0 \text{ and } t \rightarrow 0$$

follows from Proposition 1 (cf. Example 2.12 of [2] for $t = 1$). \square

In the first case of the previous corollary the dependence of good small deviation estimates and good LIL becomes transparent. The fact that we cannot specify the constants λ_1, λ_2 in (3.2) is only caused by the weak asymptotics for the small deviation estimate as we do not lose any further constants in the transfer of small deviations to the LIL. If one does not have more control on the constants λ_1, λ_2 , the understanding of the precise small time behavior of X is far from optimal as the error enters exponentially.

4. PROOFS

We start with a lemma which shows that the small deviation order is at least as large as the term induced by the variance, defined in (2.5).

Lemma 1. *Let $\varepsilon > 0$ and let X be a Lévy process with Lévy measure concentrated on $[-\varepsilon, \varepsilon]$, then*

$$\mathbb{P}(\|X\|_t \leq \varepsilon/2) \leq \exp\left(-\varepsilon^{-2} \left(\int_{-\varepsilon}^{\varepsilon} x^2 \Pi(dx) + \sigma^2\right) t/12 + 1\right), \quad \text{for } t \geq 0.$$

Proof. We proceed similarly to Lemma 4.2 in [2]. Let τ be the first exit time of X out of $[-\varepsilon, \varepsilon]$. Then, by Wald's identity,

$$\begin{aligned} 4\varepsilon^2 &\geq \limsup_{t \rightarrow \infty} \mathbb{E}[X_{t \wedge \tau}^2] \geq \limsup_{t \rightarrow \infty} \text{var}[X_{t \wedge \tau}] \\ &= \limsup_{t \rightarrow \infty} \left(\int_{-\varepsilon}^{\varepsilon} x^2 \Pi(dx) + \sigma^2\right) \mathbb{E}[t \wedge \tau] = \left(\int_{-\varepsilon}^{\varepsilon} x^2 \Pi(dx) + \sigma^2\right) \mathbb{E}[\tau]. \end{aligned}$$

Therefore,

$$\mathbb{P}\left(\tau \geq 8\varepsilon^2 / \left(\int_{-\varepsilon}^{\varepsilon} x^2 \Pi(dx) + \sigma^2\right)\right) \leq \frac{\left(\int_{-\varepsilon}^{\varepsilon} x^2 \Pi(dx) + \sigma^2\right) \mathbb{E}[\tau]}{8\varepsilon^2} \leq \frac{1}{2}.$$

Let $n := \lfloor t(\int_{-\varepsilon}^{\varepsilon} x^2 \Pi(dx) + \sigma^2)/(8\varepsilon^2) \rfloor$ and set $t_i := 8i\varepsilon^2/(\int_{-\varepsilon}^{\varepsilon} x^2 \Pi(dx) + \sigma^2)$, $i = 0, \dots, n$. Then

$$\mathbb{P}(\|X\|_t \leq \varepsilon/2) \leq \mathbb{P}\left(\forall i = 0, \dots, n-1 : \sup_{s \in [t_i, t_{i+1}]} |X_s - X_{t_i}| \leq \varepsilon\right) = \mathbb{P}(\tau \geq t_1)^n \leq 2^{-n}.$$

□

This shows that the small deviation order is always at least as large as the term induced by the truncated variance process. This fact will be needed later on.

Lemma 2. *Let F be a function that increases to infinity at zero. If for some Lévy process X for $t \leq t_0$ and $\varepsilon < \varepsilon_0$*

$$-\log \mathbb{P}(\|X\|_t \leq \varepsilon) \leq F(\varepsilon)t$$

then, for some absolute constant $c > 0$ and all $\varepsilon > 0$ small enough,

$$\varepsilon^{-2}U(\varepsilon) \leq c(F(\varepsilon) + 1).$$

Proof. We use the assumption together with the fact that if $\|X\|_t \leq \varepsilon$ then X must not have jumps larger than 2ε and the previous lemma:

$$e^{-F(\varepsilon)t} \leq \mathbb{P}(\|X\|_t \leq \varepsilon) = e^{-\bar{\Pi}(2\varepsilon)t} \mathbb{P}(\|X'\|_t \leq \varepsilon) \leq e^{-\bar{\Pi}(2\varepsilon)t} e^{-(2\varepsilon)^{-2}(\int_{-2\varepsilon}^{2\varepsilon} x^2 \Pi(dx) + \sigma^2)t/12+1},$$

where X' has Lévy measure Π restricted to $[-2\varepsilon, 2\varepsilon]$. Noting that Lemma 5.1 of [2] implies that $U(\varepsilon)/\varepsilon^2 \approx U(2\varepsilon)/(2\varepsilon)^2$, the statement of the lemma is proved. □

The lower bound in the LIL comes from the following lemma.

Lemma 3. *Let F be a function that increases to infinity at zero such that for all $t \leq t_0$ and $\varepsilon \leq \varepsilon_0$*

$$\lambda F(\varepsilon)t \leq -\log \mathbb{P}(\|X\|_t \leq \varepsilon)$$

and, for $\lambda > 0$, we set $b_\lambda(t) := F^{-1}\left(\frac{\log|\log t|}{\lambda t}\right)$. Then, for any $\lambda' < \lambda$,

$$1 \leq \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda'}(t)} \quad \text{a.s.}$$

Proof. For any $\lambda' < \lambda$, we can find $0 < r < 1$ such that $1 < \lambda r / \lambda'$. Note that

$$\sum_n \mathbb{P}(\|X\|_{r^{n+1}} \leq b_{\lambda'}(r^n)) < \infty$$

since

$$(4.1) \quad -\log \mathbb{P}(\|X\|_{r^{n+1}} \leq b_{\lambda'}(r^n)) \geq \lambda F(b_{\lambda'}(r^n)) r^n r = \lambda \frac{r}{\lambda'} \log |\log r^n| = \log n^{r\lambda/\lambda'} + \text{const.}$$

Hence, by the Borel-Cantelli lemma,

$$\{n : \|X\|_{r^{n+1}} \leq b_{\lambda'}(r^n)\}$$

is almost surely a finite set. Thus, for each path ω , we have that for any $n \geq n_0(\omega)$ and any $t \in [r^{n+1}, r^n)$

$$\frac{\|X\|_t}{b_{\lambda'}(t)} \geq \frac{\|X\|_{r^{n+1}}}{b_{\lambda'}(r^n)} \geq 1,$$

as $b_{\lambda'}$ is an increasing function. We take $\liminf_{t \rightarrow 0}$ to obtain the statement. \square

The proof of the upper bound in the LIL requires the following lemma.

Lemma 4. *Let F be a function that increases to infinity at zero such that for all $t \leq t_0$ and $\varepsilon \leq \varepsilon_0$*

$$-\log \mathbb{P}(\|X\|_t \leq \varepsilon) \leq \lambda F(\varepsilon)t$$

and, for $\lambda > 0$, set $b_\lambda(t) := F^{-1}\left(\frac{\log |\log t|}{\lambda t}\right)$. Assume that

$$(4.2) \quad \limsup_{n \rightarrow \infty} \frac{\|X\|_{(n+1)^{-(n+1)\beta}}}{b_\lambda(n^{-n\beta})} = 0 \quad \text{a.s.},$$

for all $\beta > 1$. Then, for any $\lambda' > \lambda$,

$$(4.3) \quad \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda'}(t)} \leq 1 \quad \text{a.s.}$$

Proof. For $\lambda' > \lambda$, we choose $\beta > 1$ such that $\lambda' > \lambda\beta$. First note that (4.2) implies

$$(4.4) \quad \limsup_{n \rightarrow \infty} \frac{\|X\|_{(n+1)^{-(n+1)\beta}}}{b_{\lambda'}(n^{-n\beta})} = 0 \quad \text{a.s.},$$

as $b_\lambda(t)$ is an increasing function in λ for fixed $t \geq 0$. Using the Lévy property we see the following:

$$\begin{aligned} & \sum_n \mathbb{P}\left(\sup_{(n+1)^{-(n+1)\beta} \leq t < n^{-n\beta}} |X_t - X_{(n+1)^{-(n+1)\beta}}| \leq b_{\lambda'}(n^{-n\beta})\right) \\ &= \sum_n \mathbb{P}(\|X\|_{n^{-n\beta} - (n+1)^{-(n+1)\beta}} \leq b_{\lambda'}(n^{-n\beta})) \\ &\geq \sum_n \mathbb{P}(\|X\|_{n^{-n\beta}} \leq b_{\lambda'}(n^{-n\beta})) = \infty. \end{aligned}$$

The last step follows as in (4.1) since now $\lambda\beta/\lambda' < 1$. The Borel-Cantelli lemma shows that the sequence of independent events

$$A_n = \left\{ \sup_{(n+1)^{-(n+1)\beta} \leq t < n^{-n\beta}} |X_t - X_{(n+1)^{-(n+1)\beta}}| \leq b_{\lambda'}(n^{-n\beta}) \right\}$$

satisfies $\mathbb{P}(A_n \text{ i.o.}) = 1$. To reduce to the supremum note that

$$\frac{\|X\|_{n^{-n\beta}}}{b_{\lambda'}(n^{-n\beta})} \leq \frac{\sup_{(n+1)^{-(n+1)\beta} \leq t < n^{-n\beta}} |X_t - X_{(n+1)^{-(n+1)\beta}}|}{b_{\lambda'}(n^{-n\beta})} + \frac{2\|X\|_{(n+1)^{-(n+1)\beta}}}{b_{\lambda'}(n^{-n\beta})}$$

and therefore by (4.4)

$$\liminf_{n \rightarrow \infty} \frac{\|X\|_{n^{-n\beta}}}{b_{\lambda'}(n^{-n\beta})} \leq \liminf_{n \rightarrow \infty} \frac{\sup_{(n+1)^{-(n+1)\beta} \leq t < n^{-n\beta}} |X_t - X_{(n+1)^{-(n+1)\beta}}|}{b_{\lambda'}(n^{-n\beta})} \leq 1.$$

This shows (4.3). \square

Now we are in position to prove Theorem 1. For a detailed analysis of the lim sup case, we refer to [20].

Proof of Theorem 1: The claim follows from Lemmas 3 and 4. To verify the use of Lemma 4 we still need to check that condition (4.2) holds for all $\beta > 1$.

We fix $\beta > 1$ and $\lambda'_2 > \lambda_2$. Since λ'_2 is fixed, we suppress the subscript λ'_2 in the definition of b in order to increase readability. We define the auxiliary function

$$h(t) = b(\phi(t)),$$

where $\phi(t)$ is chosen such that $\phi\left(\left(\frac{t}{t+1}\right)^{\frac{1}{\beta}}\right) = t^{\frac{1}{\beta}}$ and $\phi(0) = 0$. Note that ϕ is increasing and that $\phi(s^{-s^\beta}) = (s-1)^{-(s-1)^\beta}$. We also do not record that ϕ and h depend on β and λ'_2 .

Step 1: We show that

$$(4.5) \quad \int_0^{1/2} \bar{\Pi}(h(t)) dt < \infty.$$

First, by the definition of h and a change of variables we obtain

$$\begin{aligned} & \int_0^{1/2} \bar{\Pi}(h(t)) dt \\ &= \int_0^{C(\beta)} \bar{\Pi}(b(s^{s^{-\beta}})) \frac{d\left(\frac{s}{s+1}\right)^{\left(\frac{s+1}{s}\right)^\beta}}{ds} \\ &= \int_0^{C(\beta)} \bar{\Pi}(b(s^{s^{-\beta}})) \left(\frac{s}{s+1}\right)^{\left(\frac{s+1}{s}\right)^\beta} \left(\frac{s+1}{s}\right)^{\beta-1} s^{-2} (1 - \beta \log(1 - (s+1)^{-1})) ds, \end{aligned}$$

which can be estimated from above by

$$\begin{aligned} & C \int_0^{C(\beta)} \frac{b^2(s^{s^{-\beta}}) \bar{\Pi}(b(s^{s^{-\beta}}))}{b^2(s^{s^{-\beta}})} \left(\frac{s}{s+1}\right)^{\left(\frac{s+1}{s}\right)^\beta} s^{-1-\beta} |\log s| ds \\ &\leq C \int_0^{C(\beta)} \frac{U(b(s^{s^{-\beta}}))}{b^2(s^{s^{-\beta}})} \left(\frac{s}{s+1}\right)^{\left(\frac{s+1}{s}\right)^\beta} s^{-1-\beta} |\log s| ds \\ &\leq C' \int_0^{C(\beta)} F(b(s^{s^{-\beta}})) \left(\frac{s}{s+1}\right)^{\left(\frac{s+1}{s}\right)^\beta} s^{-1-\beta} |\log s| ds \\ &= \frac{C'}{\lambda} \int_0^{C(\beta)} \frac{\log |\log s^{s^{-\beta}}|}{s^{s^{-\beta}}} \left(\frac{s}{s+1}\right)^{\left(\frac{s+1}{s}\right)^\beta} s^{-1-\beta} |\log s| ds \\ &\leq \frac{C'}{\lambda} \int_0^{C(\beta)} s^{-1-\beta} \left(\log |\log s^{s^{-\beta}}| \right) s^{\left(\frac{s+1}{s}\right)^\beta - \frac{1}{s^\beta}} |\log s| ds < \infty, \end{aligned}$$

where we have used $x^2 \bar{\Pi}(x) \leq x^2 \bar{\Pi}(x) + \int_{-x}^x y^2 \Pi(dy) + \sigma^2 = U(x) \leq cx^2 F(x)$ for some absolute $c > 0$ by Lemma 2 and the definition of b .

Step 2: We denote by

$$(4.6) \quad A_n := \left\{ \text{there is at least one jump with modulus} > b(n^{-n^\beta}) \text{ up to time } (n+1)^{-(n+1)^\beta} \right\}$$

and show that

$$(4.7) \quad \sum_n \mathbb{P}(A_n) < \infty.$$

This comes from (4.5). Indeed, note that h inherits the monotonicity of b and ϕ and hence (4.5) implies that

$$(4.8) \quad \sum_n \left((n+1)^{-(n+1)^\beta} - (n+2)^{-(n+2)^\beta} \right) \bar{\Pi}(h((n+1)^{-(n+1)^\beta})) \leq \sum_n \int_{(n+2)^{-(n+2)^\beta}}^{(n+1)^{-(n+1)^\beta}} \bar{\Pi}(h(t)) dt < \infty.$$

Using

$$(n+1)^{-(n+1)^\beta} - (n+2)^{-(n+2)^\beta} \sim (n+1)^{-(n+1)^\beta},$$

$$b(n^{-n^\beta}) = h((n+1)^{-(n+1)^\beta}),$$

and that the sequence $(n+1)^{-(n+1)^\beta} \bar{\Pi}(h((n+1)^{-(n+1)^\beta}))$ tends to zero by (4.8), we obtain that

$$\mathbb{P}(A_n) = 1 - e^{-(n+1)^{-(n+1)^\beta} \bar{\Pi}(b(n^{-n^\beta}))} \sim (n+1)^{-(n+1)^\beta} \bar{\Pi}(h((n+1)^{-(n+1)^\beta}))$$

is summable. Therefore (4.7) is proved.

Step 3: Let us now show how to use (4.7) to deduce (4.2). Obviously, it suffices to show that

$$\limsup_{n \rightarrow \infty} \frac{\|X\|_{(n+1)^{-(n+1)^\beta}}}{b(n^{-n^\beta})} < \varepsilon \text{ a.s.},$$

for any $\varepsilon > 0$ and, hence, by the Borel-Cantelli lemma it suffices to show that

$$\sum_n \mathbb{P}(\|X\|_{(n+1)^{-(n+1)^\beta}} > \varepsilon b(n^{-n^\beta})) < \infty.$$

Separating jumps of absolute value larger or smaller than $b(n^{-n^\beta})$ and using the definition of A_n in (4.6), we obtain that

$$\begin{aligned} & \sum_n \mathbb{P}(\|X\|_{(n+1)^{-(n+1)^\beta}} > \varepsilon b(n^{-n^\beta})) \\ &= \sum_n \mathbb{P}(\|X\|_{(n+1)^{-(n+1)^\beta}} > \varepsilon b(n^{-n^\beta}); A_n^c) + \sum_n \mathbb{P}(\|X\|_{(n+1)^{-(n+1)^\beta}} > \varepsilon b(n^{-n^\beta}); A_n), \end{aligned}$$

which is bounded from above by

$$\sum_n \mathbb{P}\left(\|X\|_{(n+1)^{-(n+1)^\beta}} > \varepsilon b(n^{-n^\beta}) \mid A_n^c\right) \cdot \mathbb{P}(A_n^c) + \sum_n \mathbb{P}(A_n).$$

The second term is finite by (4.7); and the first term is bounded by

$$(4.9) \quad \sum_n \mathbb{P}\left(\|X\|_{(n+1)^{-(n+1)^\beta}} > \varepsilon b(n^{-n^\beta}) \mid A_n^c\right).$$

To estimate this sum note that conditionally on A_n^c , $X_t \stackrel{d}{=} X_t(n)$, where $X(n)$ differs from X only by removing jumps of size larger than $|b(n^{-n^\beta})|$. Clearly, by Wald's identity,

$$(4.10) \quad \text{var}(X_t(n)) = t \left(\int_{-b(n^{-n^\beta})}^{b(n^{-n^\beta})} y^2 \Pi(dy) + \sigma^2 \right) \leq tU(b(n^{-n^\beta})).$$

Note that

$$\mathbb{E}X_{(n+1)^{-(n+1)^\beta}}(n) = (n+1)^{-(n+1)^\beta} \left| \int_{|x| > b(n^{-n^\beta})} x \Pi(dx) - \gamma \right|.$$

Therefore, by assumption (2.2), taking into account that the expectation of the Lévy process $X(n)$ is linear in time,

$$\|\mathbb{E}X_{(n+1)^{-(n+1)^\beta}}(n)\|_{(n+1)^{-(n+1)^\beta}} = |\mathbb{E}X_{(n+1)^{-(n+1)^\beta}}(n)| = o(b(n^{-n^\beta})).$$

Using the previous relation (first step), Doob's martingale inequality (second step), (4.10) (third step), Lemma 2 (fourth step), and the definition of b (fifth step), we are led to the upper bound

of the term in (4.9):

$$\begin{aligned}
& \sum_n \mathbb{P} \left(\|X(n)\|_{(n+1)^{-(n+1)\beta}} > \varepsilon b(n^{-n^\beta}) \right) \\
& \leq \sum_n \mathbb{P} \left(\|X(\cdot)(n) - \mathbb{E}X(\cdot)(n)\|_{(n+1)^{-(n+1)\beta}} > \frac{1}{2} \varepsilon b(n^{-n^\beta}) \right) \\
& \leq \sum_n \frac{4\mathbb{E}|X_{(n+1)^{-(n+1)\beta}}(n) - \mathbb{E}X_{(n+1)^{-(n+1)\beta}}(n)|^2}{(\varepsilon/2)^2 b(n^{-n^\beta})^2} \\
& \leq \sum_n \frac{4(n+1)^{-(n+1)\beta} U(b(n^{-n^\beta}))}{(\varepsilon/2)^2 b(n^{-n^\beta})^2} \\
& \leq \sum_n \frac{4(n+1)^{-(n+1)\beta} C \cdot F(b(n^{-n^\beta}))}{(\varepsilon/2)^2} \\
& = \frac{C'}{\lambda \varepsilon^2} \sum_n \frac{(n+1)^{-(n+1)\beta} \log |\log n^{-n^\beta}|}{n^{-n^\beta}} < \infty,
\end{aligned}$$

where we used the definition of b in the last step. Thus, the term in (4.9) is finite, as required. \square

Proof of Corollary 1. If F is regularly varying so is b_λ , see [6], Proposition 1.5.7. Now note that if F is regularly varying with exponent $-\alpha < 0$, we have

$$\begin{aligned}
b_\lambda(t) &= F^{-1}(\log |\log t| / \lambda t) \\
&\sim \lambda^{1/\alpha} F^{-1}(\log |\log t| / t) \\
&= \lambda^{1/\alpha} b_1(t).
\end{aligned}$$

Hence, the statement of Theorem 1 reads

$$(\lambda'_1)^{1/\alpha} \leq \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_1(t)} \leq (\lambda'_2)^{1/\alpha} \quad \text{a.s.}$$

for all $\lambda'_1 < \lambda_1$ and $\lambda'_2 > \lambda_2$. Taking the limits on both sides we obtain

$$(\lambda_1)^{1/\alpha} \leq \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_1(t)} \leq (\lambda_2)^{1/\alpha} \quad \text{a.s.}$$

Applying the regular variation argument in the reverse direction yields the claim. \square

Proof of Corollary 2. This follows directly from Theorem 1. The bounds on the constants can be obtained from the absolute constants in Proposition 1. \square

Proof of Theorem 2. Lemma 3 gives the lower LIL of the theorem. Unfortunately, the arguments for the proof of Theorem 1 do not apply here. Hence, for the reverse direction we show more directly that the given norming function of the LIL implies the rate function of the small deviations. The following arguments go back to Kesten. The proof is via contradiction assuming that

$$(4.11) \quad \liminf_{t \rightarrow 0} \frac{\|X\|_t}{b_{\lambda'_2}(t)} > \frac{2}{C} + \delta$$

for some $\delta > 0$ and $\lambda'_2 > \lambda_2$. We show that under this assumption we can derive, for sufficiently large l , the estimates

$$(4.12) \quad 1 \geq \sum_{n \geq l} \mathbb{P} \left(\frac{\|X\|_{r^j - r^n}}{b_{\lambda'_2}(r^j - r^n)} > \frac{2}{C}; \text{ for all } l \leq j \leq n-1 \right) \mathbb{P}(\|X\|_{r^n} \leq b_{\lambda'_2}(r^n))$$

$$(4.13) \quad \geq \frac{1}{2} \sum_{n \geq l} \mathbb{P}(\|X\|_{r^n} \leq b_{\lambda'_2}(r^n))$$

which is a contradiction as, by the choice of $b_{\lambda'_2}$ and the small deviation rate (2.6), the sum in (4.13) is infinite. First, let us derive estimate (4.12) for which Assumption (4.11) is not needed. For any fixed integer l partitioning the probability space we obtain

$$\begin{aligned} 1 &\geq \sum_{n \geq l} \mathbb{P}(\|X\|_{r^j} > b_{\lambda'_2}(r^j) \text{ for all } l \leq j \leq n-1; \|X\|_{r^n} \leq b_{\lambda'_2}(r^n)) \\ &\geq \sum_{n \geq l} \mathbb{P}\left(\sup_{r^n \leq s < r^j} |X_s| > b_{\lambda'_2}(r^j) \text{ for all } l \leq j \leq n-1; \|X\|_{r^n} \leq b_{\lambda'_2}(r^n)\right). \end{aligned}$$

In order to employ the independence of increments of X we estimate from below by

$$\sum_{n \geq l} \mathbb{P}\left(\sup_{r^n \leq s < r^j} |X_s - X_{r^n}| > 2b_{\lambda'_2}(r^j) \text{ for all } l \leq j \leq n-1; \|X\|_{r^n} \leq b_{\lambda'_2}(r^n)\right)$$

which equals

$$\begin{aligned} &\sum_{n \geq l} \mathbb{P}(\|X\|_{r^j - r^n} > 2b_{\lambda'_2}(r^j) \text{ for all } l \leq j \leq n-1) \mathbb{P}(\|X\|_{r^n} \leq b_{\lambda'_2}(r^n)) \\ &= \sum_{n \geq l} \mathbb{P}\left(\frac{\|X\|_{r^j - r^n}}{b_{\lambda'_2}(r^j - r^n)} > 2 \frac{b_{\lambda'_2}(r^j)}{b_{\lambda'_2}(r^j - r^n)} \text{ for all } l \leq j \leq n-1\right) \mathbb{P}(\|X\|_{r^n} \leq b_{\lambda'_2}(r^n)). \end{aligned}$$

By the monotonicity of $b_{\lambda'_2}$ this yields the lower bound

$$\sum_{n \geq l} \mathbb{P}\left(\frac{\|X\|_{r^j - r^n}}{b_{\lambda'_2}(r^j - r^n)} > 2 \frac{b_{\lambda'_2}(r^j)}{b_{\lambda'_2}(r^j - r^{j+1})}; \text{ for all } l \leq j \leq n-1\right) \mathbb{P}(\|X\|_{r^n} \leq b_{\lambda'_2}(r^n)).$$

Finally, we utilize the regularity of $b_{\lambda'_2}$ from (2.8) to obtain the lower bound

$$\sum_{n \geq l} \mathbb{P}\left(\frac{\|X\|_{r^j - r^n}}{b_{\lambda'_2}(r^j - r^n)} > \frac{2}{C}; \text{ for all } l \leq j \leq n-1\right) \mathbb{P}(\|X\|_{r^n} \leq b_{\lambda'_2}(r^n)).$$

As required we derived Estimate (4.12).

Assuming (4.11) we now derive Estimate (4.13). The assumption directly shows that

$$\lim_{t \rightarrow 0} \mathbb{P}\left(\bigcap_{s \leq t} \{\|X\|_s \geq 2C^{-1}b_{\lambda'_2}(s)\}\right) = 1$$

which implies that we may choose l large enough such that

$$\mathbb{P}\left(\frac{\|X\|_{r^j - r^n}}{b_{\lambda'_2}(r^j - r^n)} > \frac{2}{C}; \text{ for all } l \leq j \leq n-1\right) \geq \mathbb{P}\left(\bigcap_{s \leq r^l} \{\|X\|_s \geq 2C^{-1}b_{\lambda'_2}(s)\}\right) \geq \frac{1}{2}.$$

Hence, we derived estimate (4.13) so that the proof is complete. \square

Proof of Corollary 3. This is completely analogous to the proof of Corollary 1. \square

Proof of Corollary 4. We use Proposition 1 and Theorem 1. In order to do so, we have to see that the term $\varepsilon u_\varepsilon$ in (2.12) has no influence on the order. We apply Lemma 3 and follow the proof of Theorem 2 with the scaling

$$t = r^n \quad \text{and} \quad \varepsilon = b(r^n)$$

and with the sequence n^{-n^β} , respectively. Therefore, it is sufficient to show that

$$\varepsilon u_\varepsilon = o(tF(\varepsilon))$$

with the above scalings of t and ε . Since $\varepsilon = b(t)$ and thus $t \sim F(\varepsilon)^{-1} \log \log F(\varepsilon)$, we need to show that

$$\varepsilon u_\varepsilon = o(\log \log F(\varepsilon)).$$

As this is precisely what we stated in condition (2.11), the proof is complete. \square

Proof of Proposition 2. As X is of bounded variation, the representation

$$X_t = A_t^1 - A_t^2 + ct$$

holds with two independent pure jump subordinators A^1, A^2 . Next, we use the simple observation

$$\frac{|X_t|}{t} \leq \frac{\|X\|_t}{t} \leq \frac{\|A^1\|_t + \|A^2\|_t + |c|t}{t} = \frac{A_t^1}{t} + \frac{A_t^2}{t} + |c|$$

to conclude the proof. The left hand side converges to $|c|$ as X has bounded variation (see Theorem 39 of [10]). Finally, the right hand side converges to $|c|$ as $|A_t^i|/t$ converge at zero almost surely to their drift (see Proposition 5 of [10]). \square

ACKNOWLEDGMENT

We thank Thomas Simon (Lille) for pointing out the reference [26] to the authors.

REFERENCES

- [1] P. Andrew. On the limiting behaviour of Lévy processes at zero. *Probability Theory and Related Fields* 140, 103–127, 2008.
- [2] F. Aurzada and S. Dereich. Small deviations of general Lévy processes. *Annals of Probability* 37, 2066–2092, 2009.
- [3] J. Bertoin. Lévy processes. Cambridge Tracts in Mathematics, 121, 1996.
- [4] J. Bertoin. On the first exit time of a completely asymmetric stable process from a finite interval. *Bull. London Math. Soc.* 28, no. 5, 514–520, 1996.
- [5] J. Bertoin, R.A. Doney, R.A. Maller. Passage of Lévy Processes across Power Law Boundaries at Small Times. *Annals of Probability* 36, 160–197, 2008.
- [6] N.H. Bingham, C.M. Goldie, and J.T. Teugels. *Regular Variation*, volume 27 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1989.
- [7] A.A. Borovkov and A.A. Mogulskii. On the probabilities of small deviations for stochastic processes. *Sib. Adv. Math.*, 1:39–63, 1991.
- [8] B. Buchmann and R.A. Maller. The small-time Chung-Wichura law for Lévy processes with non-vanishing Brownian component. doi:10.1007/s00440-009-0255-1
- [9] K.L. Chung. On the maximum partial sums of sequences of independent random variables. *Trans. Am. Math. Soc.* 64 205–233, 1948.
- [10] R. Doney. *Fluctuation theory for Lévy processes*. Springer Lecture Notes in Mathematics, 1897, 2007.
- [11] B.E. Fristedt and W.E. Pruitt. Lower functions for increasing random walks and subordinators. *Z. Wahrscheinlichkeitstheorie verw. Gebiete* 23 167–182, 1972.
- [12] W.V. Li and Q.-M. Shao. Gaussian Processes: Inequalities, small ball probabilities and applications. In: Rao, C. R. and Shanbhag, D. (eds.), *Stochastic Processes: Theory and Methods*, volume 19 of *Handbook of Statistics*, Elsevier, New York, pp. 533–597, 2001.
- [13] M.A. Lifshits. Asymptotic behavior of small ball probabilities. In: *Probab. Theory and Math. Statist. Proc. VII International Vilnius Conference*, pp. 453–468, VSP/TEV, Vilnius, 1999.
- [14] M.A. Lifshits. *Bibliography on small deviation probabilities*, available from: <http://www.proba.jussieu.fr/pageperso/smalldev/biblio.html>
- [15] M. Lifshits and T. Simon. Small deviations for fractional stable processes. *Ann. Inst. H. Poincaré Probab. Statist.* 41:725–752, 2005.
- [16] W. Linde and Z. Shi. Evaluating the small deviation probabilities for subordinated Lévy processes. *Stochastic Process. Appl.* 113:273–287, 2004.
- [17] W. Linde and P. Zipfel. Small deviation of subordinated processes over compact sets. *Probab. Math. Statist.* 28, 281–304, 2008.
- [18] A.A. Mogulskii. Small deviations in a space of trajectories. *Theor. Probab. Appl.*, 19, 726–736, 1974.
- [19] K. Sato. *Lévy processes and infinitely divisible distributions*, volume 68 in *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, Cambridge, 1999.
- [20] M. Savov. Small time two-sided LIL behavior for Lévy processes at zero. *Probability Theory and Related Fields* 144, 79–98, 2009.
- [21] T. Simon. Sur les petites déviations d’un processus de Lévy. *Potential Analysis*, 14, 155–173, 2001.
- [22] T. Simon. Small deviations in p -variation for multidimensional Lévy processes. *J. Math. Kyoto Univ.*, 43, 523–565, 2003.
- [23] E. Shmileva. Small ball probabilities for jump Lévy processes from the Wiener domain of attraction. *Statist. Probab. Lett.* 76, 1873–1881, 2006.
- [24] E. Shmileva. Shifted small deviations and Chung LIL for symmetric alpha-stable processes. arXiv:0811.2583
- [25] S.J. Taylor. Sample path properties of a transient stable process. *J. Math. Mech.*, 16, 1229–1246, 1967.
- [26] I. Wee. Lower functions for processes with stationary independent increments. *Probability Theory and Related Fields* 77, 551–566, 1988.

- [27] I. Wee. Lower functions for asymmetric Lévy processes. *Probability Theory and Related Fields* 85, 468–488, 1990.

TECHNISCHE UNIVERSITÄT BERLIN, INSTITUT FÜR MATHEMATIK, SEKR. MA 7-4, STRASSE DES 17. JUNI 136, 10623 BERLIN, GERMANY

E-mail address: `aurzada@math.tu-berlin.de`

DEPARTMENT OF STATISTICS, UNIVERSITY OF OXFORD, 1, SOUTH PARKS ROAD, OXFORD OX1 3TG, UNITED KINGDOM

E-mail address: `leif.doering@googlemail.com`

DEPARTMENT OF STATISTICS, UNIVERSITY OF OXFORD, 1, SOUTH PARKS ROAD, OXFORD OX1 3TG, UNITED KINGDOM

E-mail address: `savov@stats.ox.ac.uk`