

Existence and Structure of Stochastic Equilibria with Intertemporal Substitution[†]

BY PETER BANK AND FRANK RIEDEL[‡]

Humboldt University of Berlin

February, 2000 — this Version: December 20, 2000

Abstract

We prove existence of an Arrow–Debreu equilibrium when agents’ preferences exhibit local substitution in the sense of Hindy, Huang, and Kreps (1992). Efficient allocations and supporting price functionals are identified and characterized. Under Hindy–Huang–Kreps preferences, equilibrium price processes are semimartingales for arbitrary endowments, but in general, they do not belong to the space suggested originally by Hindy and Huang (1992) due to an additional singular local time component.

[†] Support of Deutsche Forschungsgemeinschaft through SFB 373, “Quantification and Simulation of Economic Processes”, is gratefully acknowledged.

[‡] We would like to thank Darrel Duffie for suggesting a more general approach, and Ulrich Horst for helpful discussions.

AMS 2000 subject classification. Primary 91B50; secondary 91B70

JEL subject classification. D51, D91

Key words and phrases. General Equilibrium, Local Substitution, Non–Time Additive Utility

Introduction

The most widespread utility functional used in the economics of time and uncertainty is the time-additive expected utility functional. This functional forms the basis for Merton's (1971) theory of optimal portfolio choice and Breeden's (1979) Consumption Based CAPM and the huge literature following these two fundamental articles.

The time-additive utility functional has been criticized for several deficiencies and consequently been extended in several directions. Here, we are concerned with what we consider to be the most fundamental critic made by Hindy, Huang, and Kreps (1992) (HHK henceforth) which is, at the same time, also an inspiring new approach to intertemporal decision theory.

HHK start with an economically sensible requirement for intertemporal preferences: since consumption of a good at one time is a close substitute for consumption of the same good an instant later, preferences for intertemporal consumption plans should react in a continuous way when a consumption plan is shifted slightly over time. HHK present several topologies which formalize the idea of intertemporal substitution and compute the corresponding topological duals from which price functionals should come. As a negative result, they show that the time-additive expected utility functional is not continuous with respect to those topologies — stated differently, the time-additive expected utility functional is not a good choice if one accepts the requirement of intertemporal substitution. HHK present alternative utility functionals which are indeed continuous with respect to the proposed topologies and Hindy and Huang (1992) extend this approach to the stochastic framework. They do not establish, however, the existence of an equilibrium.

In the case of certainty, the problem of existence is solved by Mas-Colell and Richard (1991) which is the most general existence theorem for exchange economies with infinite dimensional commodity space to date; in fact, the HHK paper provoked the Mas-Colell–Richard theorem.

In the general case with uncertainty, however, the Mas-Colell–Richard theorem does not apply, and existence of an equilibrium has remained an open issue. The present paper fills this gap. We prove existence of exchange equilibria for economies which include the HHK economy as a special case, and we characterize equilibrium prices.

Concerning our setup, we follow in spirit if not literally the fundamental work of Hindy, Huang, and Kreps (1992).

Consumption patterns are described in cumulative form, i.e., as rightcontinuous, increasing processes on $[0, T]$. Of course, these patterns also have to be adapted to the information flow in the economy. The consumption space is thus the positive cone in the space of all optional random measures with integrable total variation.

To capture the substitutability of consumption over time, we endow the consumption space with the topology of weak convergence in probability plus L^1 -convergence of total masses. The norm(s) used by Hindy and Huang (1992) induce the same topology on the consumption space.

Agents are characterized by strictly monotone, strictly concave utility functionals which are continuous in our topology. Moreover, we assume that subgradients exist for the utility functionals and that these subgradients are bounded and bounded away from zero on the set of feasible allocations. This smoothness assumption plays here the same role as (and is somewhat stronger than) uniform properness. The utility functionals of Hindy–Huang–Kreps and generalizations in the spirit of stochastic differential utility satisfy these assumptions.

One can show that the setting satisfies all conditions for the Mas–Colell–Richard theorem except one: the topological dual is not a lattice, or stated differently, the maximum of two price candidates does not necessarily induce a *continuous* linear functional in the Hindy and Huang (1992) model. The reason for this is inherently stochastic: local time terms appearing in the Doob–Meyer decomposition of price processes. Hindy and Huang (1992) show that the topological dual of the commodity space consists of semimartingales with absolutely continuous compensator. Now the maximum of two semimartingales is again a semimartingale; however, the compensator of such a maximum involves local time terms which in general (and in particular in the Brownian models mostly used in finance) can be singular with respect to Lebesgue measure. Thus, the compensator is not absolutely continuous in general and the topological dual is not a lattice. Hence, the Mas–Colell–Richard theorem does not apply, and a different approach is required.

Our approach keeps as much as possible from the well established tools of the literature. Indeed, as usual in the context of infinite dimensional commodity spaces with finitely many agents, the Negishi–method is the basis for the proof of existence. As in Mas-Colell and Richard (1991), a disaggregated version of the Negishi approach is used in the sense that we consider the utility gradient of every individual agent at some Pareto optimum and take as price candidate a weighted maximum of these gradients. This is where the lattice structure plays a role in the abstract Mas–Colell–Richard theorem and where, therefore, we must deviate.

The major deviation from Mas-Colell and Richard (1991) is that, since we cannot work with the topological dual of Hindy and Huang (1992), we define the topology only on the positive cone of the commodity space. Several extensions of the topology to the whole space are possible, but we do not need to specify one. The topological dual plays no role in our argument.

Instead, we employ continuity of the price candidates only on the order ideal which is generated by aggregate endowment. This order ideal can be identified as some L^∞ -space and thus, every bounded process induces a continuous linear functional. This makes it possible to prove existence of equilibria in the economy restricted to the order ideal. The same idea has been used by Duffie and Zame (1989) to establish existence of a CCAPM equilibrium with time-additive preferences.

In a final step, we show that the equilibrium of the restricted economy is also an equilibrium of the original economy. In order to perform this step, we show that supporting price functionals can be represented as weighted maxima of utility gradients evaluated at the associated optimal allocation. This gives us the required extension of the equilibrium price functional from the order ideal to the consumption space.

As explained above, the equilibrium price functional does not necessarily belong to the topological dual of Hindy and Huang (1992), i.e. there may be points of discontinuity in the *commodity space*. Nevertheless, one may ask whether the functional is continuous on the *consumption space*, i.e. the positive cone of the commodity space. We show that the answer to this question is affirmative, under the additional assumptions that the information flow in the economy is quasi-leftcontinuous and that utility gradients are semimartingales with a continuous compensator. Hence, for HHK utilities, for example, there exists an equilibrium with a price functional which is continuous on the consumption space albeit not on the whole commodity space. For the purposes of agents, this is the relevant kind of continuity since they are only interested in positive consumption bundles.

If utility gradients are semimartingales, so are equilibrium prices as they take the form of a weighted maximum of such gradients. This is notable since semimartingale prices can be ensured in the time-additive model only by assuming that the aggregate endowment rate follows a semimartingale. Here, price processes have this particularly nice structure for *any* kind of endowment stream.

In contrast to a conjecture in Hindy and Huang (1992), however, it is not possible to ensure existence of an interest rate in general. Technically, this follows from an application of the Ito–Tanaka formula to the mentioned maximum representation of equilibrium prices. It turns out that the predictable part of equilibrium prices typically has a singular local time component, even when the basic utility gradients are compensated by absolutely continuous processes. The economic reason for this singularity is that agents refrain from consuming totally when prices are too high as compared to their marginal utility of consumption, and they only resume consumption after a decline of prices. Hence, it may happen that the identity of the agent whose gradient sets the price changes when one agent stops consuming and another one restarts. If prices fluctuate in a diffusion-like manner, such periods of changes may be arbitrarily small, leading to a singular component in the evolution of equilibrium prices. Essentially the same singularity effect can also be observed in time-additive settings when marginal utility at zero is finite as has been shown by Karatzas, Lehoczky, and Shreve (1991). A difference to the time-additive case with finite marginal utility at zero is that an interest rate does exist as long as there is one agent who consumes all the time because then the price process is identical to this agent's utility gradient.

The paper is organized as follows. Section 1 describes the economy under uncertainty, the consumption space and agents' utility functionals. Section 2 analyzes efficient allocations and their supporting prices. Section 3 establishes existence of general equilibrium for this economy. Finally, the continuity properties of equilibrium price functionals are studied in Section 4.

1 The Economy

We consider a stochastic pure exchange economy with a finite set of agents \mathcal{I} . Uncertainty is modeled through a filtered probability space $(\Omega, \mathcal{F}_T, (\mathcal{F}_t, 0 \leq t \leq T), \mathbb{P})$ satisfying the usual conditions of right continuity and completeness; \mathcal{F}_0 is \mathbb{P} -a.s. trivial.

1.1 Consumption Plans and Price Functionals

Under certainty, cumulative consumption is represented by an increasing, rightcontinuous function in

$$\mathcal{M}_+ \triangleq \{F : [0, T] \rightarrow [0, +\infty) \mid F \text{ is rightcontinuous and increasing}\}.$$

Since small shifts over time affect an agent's appreciation of a consumption plan only slightly, the economically appropriate notion of closeness between consumption plans is the Prohorov distance

$$d_{\mathcal{M}_+}(F, F') \triangleq \inf\{\epsilon > 0 \mid F(t - \epsilon) - \epsilon \leq F'(t) \leq F(t + \epsilon) + \epsilon \text{ for all } t \in [0, T]\}.$$

This metric induces the weak topology on \mathcal{M}_+ . In so far we follow, in spirit if not literally, the fundamental work of Hindy, Huang, and Kreps (1992).

Under uncertainty, an agent chooses a random element $C(\omega) \in \mathcal{M}_+$ which has to conform to the available information in the sense that $(C(t))_{0 \leq t \leq T}$ is an adapted process. We assume in addition that expected total consumption remains finite and, thus, the consumption space is

$$\mathcal{X}_+ \triangleq \{C : (\Omega, \mathcal{F}_T) \rightarrow \mathcal{M}_+ \mid (C(t))_{0 \leq t \leq T} \text{ is adapted with } \mathbb{E}C(T) < +\infty\}.$$

A natural extension of the Prohorov distance to the uncertain framework is given by

$$d_{\mathcal{X}_+}(C, C') \triangleq \mathbb{E} [d_{\mathcal{M}_+}(C, C') \wedge 1] + \mathbb{E} |C(T) - C'(T)|.$$

This metric endows \mathcal{X}_+ with the topology of weak convergence in probability plus L^1 -convergence of total masses.

The consumption space \mathcal{X}_+ has a natural ordering \preceq given by

$$C \preceq C' \Leftrightarrow C' - C \in \mathcal{X}_+.$$

Remark 1.1 *Note that \preceq does not denote a preference relation. Throughout this paper, agents' preferences will be described by utility functionals. Hence, no confusion should arise.*

Every bounded measurable process $\psi : \Omega \times [0, T] \rightarrow \mathbb{R}$ gives rise to a (not necessarily continuous) linear functional $\langle \psi, \cdot \rangle$ on \mathcal{X}_+ via

$$\langle \psi, X \rangle \triangleq \mathbb{E} \int_0^T \psi(t) dX(t) \quad (X \in \mathcal{X}_+).$$

If, in addition, ψ is nonnegative and optional, we call it a price process, and we call $\langle \psi, \cdot \rangle$ a price functional on \mathcal{X}_+ .

Remark 1.2 *Note that we do not assume continuity of price functionals a priori as is usually done in equilibrium theory. In fact, we will first identify an equilibrium price functional in the much larger space of all linear functionals and establish a posteriori its continuity under appropriate conditions; see Theorem 4.1.*

1.2 Primitives of the Economy

Each agent $i \in \mathcal{I}$ is endowed with some cumulative income stream $E_i \in \mathcal{X}_+$. To avoid trivial cases, we assume $E_i \neq 0$. Aggregate endowment is $E \triangleq \sum_i E_i$.

Assumption 1 *Agent i 's preferences are described by a utility functional $V_i : \mathcal{X}_+ \rightarrow \mathbb{R}$ with the following properties:*

- (i) V_i is continuous with respect to $d_{\mathcal{X}_+}$, strictly concave and strictly increasing with respect to \preceq ,
- (ii) for every $C \in \mathcal{X}_+$ there exists a bounded optional process $\nabla V_i(C)$ with the subgradient property

$$V_i(C') - V_i(C) \leq \langle \nabla V_i(C), C' - C \rangle \quad (C' \in \mathcal{X}_+).$$

These subgradients are continuous in the sense that, for any two $C, C' \in \mathcal{X}_+$,

$$\lim_{\epsilon \downarrow 0} \langle \nabla V_i(\epsilon C' + (1 - \epsilon)C), C' - C \rangle = \langle \nabla V_i(C), C' - C \rangle.$$

Hence, we assume essentially convexity of preferences and a sufficient degree of smoothness. In addition, we require the following technical

Assumption 2 *Subgradients are uniformly bounded from above and bounded away from zero in the sense that there are nonnegative, optional processes $b, B \in L_+^1(\mathbb{P} \otimes dE) \setminus \{0\}$ such that*

$$b \leq \nabla V_i(C) \leq B \quad \mathbb{P} \otimes dE\text{-a.e.}$$

for all $C \in \mathcal{X}_+$ with $C \preceq E$.

1.3 Examples

A class of utility functionals which satisfy Assumptions 1 and 2, and which in fact motivated this work are

Hindy–Huang–Kreps Preferences. These preferences are given by a utility functional of the form

$$V(C) \triangleq \mathbb{E} \int_0^T u(t, Y(C)(t)) dt \quad (C \in \mathcal{X}_+)$$

where $u : [0, T] \times \mathbb{R}_+ \rightarrow \mathbb{R}$ denotes a continuous felicity function which is strictly concave, twice continuously differentiable and increasing in its second argument. The quantity

$$Y(C)(t) \triangleq \eta e^{-\beta t} + \int_0^t \beta e^{-\beta(t-s)} dC(s)$$

describes the investor's level of satisfaction obtained from his consumption up to time $t \in [0, T]$; $\eta > 0$ is his initial level of satisfaction, and $\beta > 0$ measures how fast satisfaction decays.

The subgradient ∇V at a given consumption plan $C \in \mathcal{X}_+$ is given by the optional version of

$$\nabla V(C)(t) \triangleq \mathbb{E} \left[\int_t^T \partial_y u(s, Y(C)(s)) \beta e^{-\beta(s-t)} ds \middle| \mathcal{F}_t \right] \quad (0 \leq t \leq T)$$

as has already been shown by Duffie and Skiadas (1994); Bank and Riedel (2000) prove continuity of ∇V in the sense of the second part of Assumption 1 (ii). The gradient bounds of Assumption 2 can be chosen as

$$b \triangleq \min_i \nabla V_i(E) \quad \text{and} \quad B \triangleq \max_i \nabla V_i(0).$$

Remark 1.3 *The above Hindy–Huang–Kreps utility is strictly increasing and strictly concave only on the slightly smaller consumption space $\{C \in \mathcal{X}_+ \mid \Delta C(T) = 0\}$, since consumption made at time $t = T$ obviously does not contribute to agent's utility. However, this minor deviation from Assumption 1 does not impose any problems if we assume $\Delta E(T) = 0$.*

The generalization of Hindy, Huang, and Zhu (1997) based on two levels of satisfaction is a further example.

Stochastic Differential Utility. Under suitable assumptions, also the stochastic differential utility version of Hindy–Huang–Kreps preferences forms an example for our setup. Indeed, let $f : [0, T] \times \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function with the following properties:

- (i) for $s \in [0, T]$, $f(s, \cdot, \cdot)$ is strictly increasing, strictly concave, and continuously differentiable,
- (ii) for $s \in [0, T]$ and $y \in \mathbb{R}_+$, $f(s, y, \cdot)$ is Lipschitz,

(iii) for $(s, y, u) \in [0, T] \times \mathbb{R}_+ \times \mathbb{R}$, the derivative with respect to y satisfies

$$\partial_y f(s, y, u) \leq K(1 + y)$$

for some constant $K > 0$.

Using similar arguments as in Duffie and Epstein (1992), one obtains for any $C \in \mathcal{X}_+$ a unique optional process $U(C)$ such that

$$U(C)(t) = \mathbb{E} \left[\int_t^T f(s, Y(C)(s), U(C)(s)) \beta e^{-\beta(s-t)} ds \middle| \mathcal{F}_t \right] \quad (0 \leq t \leq T)$$

with $Y(C)$ as above. The utility functional given by $V(C) \triangleq U(C)(0)$ satisfies then Assumption 1; compare Duffie and Skiadas (1994) and Duffie and Epstein (1992). The subgradient takes the form

$$\nabla V(C)(t) = \mathbb{E} \left[\int_t^T e^{\delta(s)} \partial_y f(s, Y(C)(s), U(C)(s)) \beta e^{-\beta(s-t)} ds \middle| \mathcal{F}_t \right]$$

where $\delta(t) \triangleq \int_0^t \partial_u f(s, Y(C)(s), U(C)(s)) ds$ ($0 \leq t \leq T$).

Remark 1.4 *Standard time-additive utility functionals like*

$$V(C) = \mathbb{E} \int_0^T u(t, c(t)) dt$$

for absolutely continuous $C(t) = \int_0^t c(s) ds$ are not continuous with respect to $d_{\mathcal{X}_+}$ and are thus **no** examples for our setup; compare Hindy, Huang, and Kreps (1992).

1.4 Equilibrium

An *allocation* is a vector $(C_i)_{i \in \mathcal{I}} \in \mathcal{X}_+^{\mathcal{I}}$. It is *feasible* if $\sum_i C_i \preceq E$. The set of feasible allocations will be denoted by \mathcal{Z} .

An (Arrow–Debreu) *equilibrium* consists of a feasible allocation $(C_i^*)_{i \in \mathcal{I}} \in \mathcal{Z}$ and a price process ψ^* such that, for any $i \in \mathcal{I}$, the consumption plan C_i^* maximizes agent i 's utility over all C_i satisfying the budget–constraint $\langle \psi^*, C_i \rangle \leq \langle \psi^*, E_i \rangle$.

The main theorem of this paper is

Theorem 1.5 *Under Assumptions 1 and 2, an equilibrium exists.*

We have given an informal sketch of the method of proof in the introduction. In the following, we focus on the technical issues involved.

In Section 2.1 we start with an analysis of the set of Pareto optima parametrized by the set of utility weights $\Lambda = \{ \lambda \in \mathbb{R}_+^{\mathcal{I}} \mid \sum_i \lambda_i = 1 \}$. Technically, it is important that the Pareto optima depend continuously on the utility weights in some topology. Such a continuity is established for the weak* topology on the order ideal in Lemma 2.2.

Let us also mention that Kabanov's version of Komlós' theorem (Kabanov (1999), Lemma 3.5; Komlós (1967)) plays an important role in the proofs. It is used to establish existence of efficient allocations, and it is also useful in showing that the efficient allocation depends continuously on agents' weights.

In Section 2.2, we characterize the price functionals which support the efficient allocations on the order ideal. The corresponding price processes are identified as weighted maxima of agents' utility gradients at the efficient allocation. This particular structure allows us to extend the supporting price functionals from the order ideal to the whole consumption space later on.

Finally, we define in Section 3 the usual excess utility correspondence. Upper hemi-continuity of this correspondence follows from a classical argument due to Bewley (1969), and Kakutani's fixed point theorem yields existence of an equilibrium for the economy restricted to the order ideal. The characterization of supporting prices as weighted maxima of utility gradients allows us to extend this equilibrium to an equilibrium of the original economy.

2 Efficient Allocations and Supporting Prices

The first step in our program is to characterize efficient allocations. To this end, we prove a version of the Kuhn–Tucker theorem for the welfare maximization problem. As usual, if agents consume, their utility gradients are equalized in an efficient allocation. The common value for the utility gradients is the Lagrange multiplier for this problem. At the same time, it gives rise to a price functional which supports the efficient allocation in the sense of the Second Welfare Theorem. Of course, there may be other functionals with the same supporting property. However, as we shall see, they all share the same structure.

2.1 The Social Welfare–Problem

Let us introduce the set of normalized weights

$$\Lambda \triangleq \left\{ \lambda \in \mathbb{R}_+^{\mathcal{I}} \mid \sum_i \lambda_i = 1 \right\}.$$

An allocation $(C_i)_{i \in \mathcal{I}}$ is called λ -efficient for agents' weights $\lambda \in \Lambda$ if it maximizes the social welfare $\sum_i \lambda_i V_i(C_i)$ subject to the feasibility constraint

$$\sum_i C_i \preceq E.$$

The characterization of efficient allocations is achieved by the following Kuhn–Tucker–like result.

Lemma 2.1 *For any $\lambda \in \Lambda$, there exists a unique λ -efficient allocation $(C_i^\lambda)_{i \in \mathcal{I}} \in \mathcal{Z}$. It is characterized by the joint validity of the following properties (i)–(iii) for some nonnegative, optional random variable ψ :*

- (i) $\sum_i C_i^\lambda = E$,
- (ii) $\lambda_i \nabla V_i(C_i^\lambda) \leq \psi$,
- (iii) $\langle \psi - \lambda_i \nabla V_i(C_i^\lambda), C_i^\lambda \rangle = 0$ for every $i \in \mathcal{I}$.

The random variable ψ plays the role of a Lagrange multiplier for the problem of maximizing social welfare. By the flat-off condition (iii), it is uniquely determined $\mathbb{P} \otimes dE$ -almost everywhere as

$$\psi = \max_i \{\lambda_i \nabla V_i(C_i^\lambda)\}.$$

PROOF : Uniqueness of the λ -efficient allocation follows as usual from the strict concavity of the utility functionals V_i ($i \in \mathcal{I}$) by considering a convex combination of two λ -efficient allocations.

To prove existence, we choose a welfare-maximizing sequence of feasible allocations $((C_i^n)_{i \in \mathcal{I}}, n = 1, 2, \dots)$ such that

$$\sum_i \lambda_i V_i(C_i^n) \rightarrow \sup_{(C_i) \in \mathcal{Z}} \sum_i \lambda_i V_i(C_i).$$

By feasibility, each sequence $(C_i^n(T), n = 1, 2, \dots)$ ($i \in \mathcal{I}$) is bounded in $L^1(\mathbb{P})$. Hence, we may use Kabanov's version of Komlós' theorem (Kabanov (1999), Lemma 3.5; Komlós (1967)) to obtain existence of a subsequence, again denoted by n , such that each sequence $(C_i^n, n = 1, 2, \dots)$ ($i \in \mathcal{I}$) is almost surely weakly Cesaro convergent to some C_i^* ($i \in \mathcal{I}$), i.e., we have almost surely

$$\tilde{C}_i^n(t) \triangleq \frac{1}{n} \sum_{k=1}^n C_i^k(t) \rightarrow C_i^*(t)$$

for $t = T$ and also for every point of continuity t of C_i^* . The above convergence shows that $(\tilde{C}_i^n)_{i \in \mathcal{I}}$ also is a feasible allocation. Moreover, it implies $d_{\mathcal{X}_+}(\tilde{C}_i^n, C_i^*) \rightarrow 0$, as, in addition to the above weak convergence, $(\tilde{C}_i^n(T), n = 1, 2, \dots)$ is dominated since

$$0 \leq \tilde{C}_i^n(T) \leq E(T)$$

by feasibility. Finally, also $((\tilde{C}_i^n)_{i \in \mathcal{I}}, n = 1, 2, \dots)$ is a maximizing sequence of feasible allocations due to concavity of social welfare. As $\tilde{C}_i^n \rightarrow C_i^*$ in $(\mathcal{X}_+, d_{\mathcal{X}_+})$ for every $i \in \mathcal{I}$, this implies λ -efficiency of $(C_i^*)_{i \in \mathcal{I}}$ by continuity of preferences (Assumption 1 (i)).

In order to prove the asserted characterization of efficient allocations, we proceed in three steps.

1. We start with sufficiency of (i)–(iii). Let $(C_i^\lambda)_{i \in \mathcal{I}}$ denote the λ -efficient allocation and let $(C_i)_{i \in \mathcal{I}} \in \mathcal{Z}$ be another feasible allocation. Due to the subgradient estimate of Assumption 1 (ii), we have

$$\sum_i \lambda_i \{V_i(C_i^\lambda) - V_i(C_i)\} \geq \sum_i \lambda_i \langle \nabla V_i(C_i^\lambda), C_i^\lambda - C_i \rangle$$

which by (ii) and (iii) is

$$\geq \left\langle \psi, \sum_i C_i^\lambda - \sum_i C_i \right\rangle.$$

As ψ is nonnegative, this yields

$$\sum_i \lambda_i \{V_i(C_i^\lambda) - V_i(C_i)\} \geq \left\langle \psi, E - \sum_i C_i \right\rangle \geq 0$$

by condition (i) and feasibility of $(C_i)_{i \in \mathcal{I}}$. Hence an allocation $(C_i)_{i \in \mathcal{I}}$ with (i)–(iii) indeed attains maximal social welfare among all feasible allocations, given agents' weights λ .

2. Necessity of condition (i) follows immediately from strict monotonicity of preferences. To prove that conditions (ii) and (iii) hold true for some Lagrange parameter ψ , consider another feasible allocation $(C_i)_{i \in \mathcal{I}} \in \mathcal{Z}$. For $\epsilon \in [0, 1]$, let $C_i^\epsilon \triangleq \epsilon C_i + (1 - \epsilon)C_i^\lambda$ ($i \in \mathcal{I}$). Since every allocation $(C_i^\epsilon)_{i \in \mathcal{I}}$ is feasible, λ -efficiency of $(C_i^\lambda)_{i \in \mathcal{I}}$ yields, for $\epsilon \in (0, 1]$,

$$\begin{aligned} 0 &\geq \frac{1}{\epsilon} \sum_i \lambda_i \{V_i(C_i^\epsilon) - V_i(C_i^\lambda)\} \\ &\geq \frac{1}{\epsilon} \sum_i \lambda_i \langle \nabla V_i(C_i^\epsilon), C_i^\epsilon - C_i^\lambda \rangle = \sum_i \lambda_i \langle \nabla V_i(C_i^\epsilon), C_i - C_i^\lambda \rangle. \end{aligned}$$

Using continuity of subgradients (Assumption 1 (ii)), we may let $\epsilon \downarrow 0$ in the above estimate to deduce

$$\sum_i \langle \phi_i, C_i \rangle \leq \sum_i \langle \phi_i, C_i^\lambda \rangle$$

where $\phi_i \triangleq \lambda_i \nabla V_i(C_i^\lambda)$ ($i \in \mathcal{I}$).

We see that $(C_i^\lambda)_{i \in \mathcal{I}}$ also solves the linear problem to maximize $\sum_i \langle \phi_i, C_i \rangle$ over all feasible allocations. In Step 3 below, we show that every solution $(C_i^*)_{i \in \mathcal{I}}$ of this problem satisfies $\langle \psi - \phi_i, C_i^* \rangle = 0$ for every $i \in \mathcal{I}$ where $\psi \triangleq \max_i \phi_i$. For $C^* = C^\lambda$, we find that, with this choice of ψ , conditions (ii) and (iii) in fact hold true, too.

3. Let $(C_i^*)_{i \in \mathcal{I}} \in \mathcal{Z}$ be a feasible allocation such that

$$\sum_i \langle \phi_i, C_i \rangle \leq \sum_i \langle \phi_i, C_i^* \rangle$$

for every other feasible allocation $(C_i)_{i \in \mathcal{I}} \in \mathcal{Z}$.

Consider the allocation defined by the optional random measures

$$dC_i(t) \triangleq n(t)^{-1} \mathbf{1}_{\{\phi_i(t) = \psi(t)\}} dE(t) \quad (i \in \mathcal{I})$$

with $\psi(t) = \max_i \phi_i(t)$ and where $n(t) \triangleq \sum_i 1_{\{\phi_i(t)=\psi(t)\}}$ denotes the number of indices i realizing the max in $\max_i \phi_i(t)$ ($0 \leq t \leq T$). Clearly, $(C_i)_{i \in \mathcal{I}}$ is feasible and satisfies

$$\sum_i \langle \phi_i, C_i \rangle = \langle \psi, E \rangle .$$

Therefore, $\sum_i \langle \phi_i, C_i^* \rangle$ cannot be less than $\langle \psi, E \rangle$. On the other hand, since $\psi \geq \phi_i \geq 0$ for each $i \in \mathcal{I}$ and as $(C_i^*)_{i \in \mathcal{I}}$ is feasible, $\sum_i \langle \phi_i, C_i^* \rangle$ cannot be greater than $\langle \psi, E \rangle$. Thus, we must in fact have

$$\sum_i \langle \phi_i, C_i^* \rangle = \langle \psi, E \rangle$$

which can hold true only if $\langle \psi - \phi_i, C_i^* \rangle = 0$ for every $i \in \mathcal{I}$.

□

The dependence of efficient allocations and induced expected utilities on agents' weights will be described in the following Lemma 2.2 and its Corollary 2.3. The latter result will be a corner stone for our proof of Theorem 1.5.

Lemma 2.2 *The mapping $\lambda \mapsto C^\lambda$ is weakly continuous in the sense that*

$$\lim_{\lambda^n \rightarrow \lambda^0} \langle \psi, C_i^{\lambda^n} \rangle = \langle \psi, C_i^{\lambda^0} \rangle \quad (i \in \mathcal{I})$$

for every $\psi \in L^1(\mathbb{P} \otimes dE)$.

PROOF : Let λ^n ($n = 1, 2, \dots$) tend to λ^0 in Λ . Put $C^n \triangleq C^{\lambda^n}$ and consider the densities

$$D_i^n \triangleq \frac{dC_i^n}{dE} \quad (n = 0, 1, \dots, i \in \mathcal{I}).$$

Due to feasibility of efficient allocations, these densities have optional versions taking values in $[0, 1]$. Now, writing

$$\langle \psi, C_i^n \rangle = \mathbb{E} \int_0^T \psi(t) D_i^n(t) dE(t) \quad (n = 0, 1, \dots, i \in \mathcal{I})$$

for $\psi \in L^1(\mathbb{P} \otimes dE)$, we see that our assertion is equivalent to the assertion that, for every $i \in \mathcal{I}$ the densities D_i^n ($n = 1, 2, \dots$) converge to D_i^0 in the weak*-topology $\sigma(L^\infty(\mathbb{P} \otimes dE), L^1(\mathbb{P} \otimes dE))$. As the unit ball in $L^\infty(\mathbb{P} \otimes dE)$ is sequentially compact with respect to this topology, this convergence will be proved once we know that, for each $i \in \mathcal{I}$, all weak*-convergent subsequences of $(D_i^n, n = 1, 2, \dots)$ have the same limit D_i^0 .

To prove this, we slightly abuse notation and suppose that $((D_i^n)_{i \in \mathcal{I}}, n = 1, 2, \dots)$ is a subsequence such that each component $(D_i^n, n = 1, 2, \dots)$ is weak*-convergent to some $D_i^\infty \in L^\infty(\mathbb{P} \otimes dE)$ ($i \in \mathcal{I}$). We then have to show that $D_i^\infty = D_i^0$ for every $i \in \mathcal{I}$.

For this, we note first that, by feasibility, we have $\mathbb{E}C_i^n(T) \leq \mathbb{E}E(T)$ for any $n = 1, 2, \dots$ and every $i \in \mathcal{I}$. Hence, by Kabanov's version of Komlós' theorem, there is a subsequence, again denoted by $((C_i^n)_{i \in \mathcal{I}}, n = 1, 2, \dots)$, such that, for any $i \in \mathcal{I}$,

$$\tilde{C}_i^n \triangleq \frac{1}{n} \sum_{k=1}^n C_i^k$$

almost surely converges weakly to some $C_i^* \in \mathcal{X}_+$ in the sense that

$$d_{\mathcal{M}_+}(\tilde{C}_i^n, C_i^*) \rightarrow 0 \quad (n \uparrow +\infty).$$

By dominated convergence, we also have $\tilde{C}_i^n(T) \rightarrow C_i^*(T)$ in $L^1(\mathbb{P})$ and, therefore, even $d_{\mathcal{X}_+}(\tilde{C}_i^n, C_i^*) \rightarrow 0$ for every $i \in \mathcal{I}$. Clearly, the densities

$$\tilde{D}_i^n \triangleq \frac{d\tilde{C}_i^n}{dE} = \frac{1}{n} \sum_{k=1}^n D_i^k \quad (n = 1, 2, \dots)$$

inherit weak*-convergence to D_i^∞ from $(D_i^n, n = 1, 2, \dots)$. Therefore, each C_i^* is almost surely absolutely continuous with respect to E with density D_i^∞ . Hence, in order to conclude our claim, we only need to prove that $(C_i^*)_{i \in \mathcal{I}}$ is the unique (!) efficient allocation for agents' weights λ^0 .

To this end, consider any other feasible allocation $C = (C_i)_{i \in \mathcal{I}} \in \mathcal{Z}$. By continuity of preferences, we have

$$\sum_i \lambda_i^0 V_i(C_i^*) = \lim_n \sum_i \lambda_i^0 V_i(\tilde{C}_i^n)$$

which, due to the concavity of every $V_i(\cdot)$ ($i \in \mathcal{I}$), is

$$\begin{aligned} &\geq \limsup_n \frac{1}{n} \sum_{k=1}^n \sum_i \lambda_i^0 V_i(C_i^k) \\ &= \limsup_n \frac{1}{n} \sum_{k=1}^n \sum_i \{\lambda_i^k V_i(C_i^k) + R_i^k\} \end{aligned}$$

for $R_i^k \triangleq (\lambda_i^0 - \lambda_i^k) V_i(C_i^k)$ ($k = 1, 2, \dots, i \in \mathcal{I}$). This term tends to zero for $k \uparrow \infty$ as $V_i(C_i^k) \in [V_i(0), V_i(E)]$ is uniformly bounded and $\lambda^k \rightarrow \lambda^0$ ($k \uparrow \infty$). Hence, we obtain

$$\sum_i \lambda_i^0 V_i(C_i^*) \geq \limsup_n \frac{1}{n} \sum_{k=1}^n \sum_i \lambda_i^k V_i(C_i^k)$$

By λ^k -efficiency of allocation C^k ($k = 1, 2, \dots$), this is in turn

$$\begin{aligned} &\geq \limsup_n \frac{1}{n} \sum_{k=1}^n \sum_i \lambda_i^k V_i(C_i) \\ &= \limsup_n \sum_i \left(\frac{1}{n} \sum_{k=1}^n \lambda_i^k \right) V_i(C_i) \\ &= \sum_i \lambda_i^0 V_i(C_i) \end{aligned}$$

where the last equality is due to the convergence $\lambda^n \rightarrow \lambda^0$. Hence, we have shown

$$\sum_i \lambda_i^0 V_i(C_i^*) \geq \sum_i \lambda_i^0 V_i(C_i)$$

for every feasible allocation $(C_i)_{i \in \mathcal{I}}$. Since, in addition, $(C_i^*)_{i \in \mathcal{I}}$ is feasible, it must coincide with the unique λ^0 -efficient allocation $(C_i^0)_{i \in \mathcal{I}}$ and we are done. \square

Let us note the following crucial

Corollary 2.3 *Every mapping $\lambda \mapsto V_i(C_i^\lambda)$ ($i \in \mathcal{I}$) is upper-semicontinuous, i.e.,*

$$\limsup_{\lambda^n \rightarrow \lambda^0} V_i(C_i^{\lambda^n}) \leq V_i(C_i^{\lambda^0}).$$

PROOF : By concavity we have

$$V_i(C_i^{\lambda^n}) - V_i(C_i^{\lambda^0}) \leq \left\langle \nabla V_i(C_i^{\lambda^0}), C_i^{\lambda^n} - C_i^{\lambda^0} \right\rangle.$$

By Lemma 2.2, the last term tends to zero if $\lambda^n \rightarrow \lambda^0$ in Λ and we are done. \square

2.2 Supporting Prices on the Order Ideal

For some of the following more technical arguments, it will be convenient to work with the auxiliary consumption space given by the order ideal

$$\tilde{\mathcal{X}}_+ = \tilde{\mathcal{X}}_+(E) \triangleq \left\{ X \in \mathcal{X}_+ \mid \frac{dX}{dE} \text{ exists } \mathbb{P}\text{-a.s. and is } \mathbb{P} \otimes dE\text{-essentially bounded} \right\}.$$

For a general discussion of the order ideal in general equilibrium theory, we refer the reader to Mas-Colell and Zame (1991).

Remark 2.4 *The space $\tilde{\mathcal{X}}_+$ can be identified with $L_+^\infty(\Omega \times [0, T], \mathcal{O}, \mathbb{P} \otimes dE)$, the set of all nonnegative, optional processes which are $\mathbb{P} \otimes dE$ -essentially bounded. Clearly, any feasible allocation is contained in $\tilde{\mathcal{X}}_+^T$.*

By Lemma 2.1, we may associate to every efficient allocation $(C_i^\lambda)_{i \in \mathcal{I}}$ ($\lambda \in \Lambda$) the Lagrange multiplier

$$\psi_\lambda \triangleq \max_i \{ \lambda_i \nabla V_i(C_i^\lambda) \}.$$

We will have a lot more to say about the structure of these multipliers in Section 4. For the moment, let us content ourselves by noting that each ψ_λ can be viewed as a nonnegative, optional random variable in $L^1(\mathbb{P} \otimes dE)$. Hence, each of these Lagrange multipliers gives rise to a price density on $\tilde{\mathcal{X}}_+$. Considering ψ_λ as a price density is also sustained by the fact that it supports its associated efficient allocation $(C_i^\lambda)_{i \in \mathcal{I}}$. This will be proved in Lemma 2.6 below.

Beforehand, let us recall that a price density $\psi^* \in L^1(\mathbb{P} \otimes dE)$ supports an allocation $(C_i^*)_{i \in \mathcal{I}}$ with $\sum_i C_i^* = E$, if it is non-zero and if any preferred allocation $(C_i)_{i \in \mathcal{I}} \in \tilde{\mathcal{X}}_+^{\mathcal{I}}$ has a higher ‘price’ under ψ^* than $(C_i^*)_{i \in \mathcal{I}}$. More precisely, we say ψ^* supports $(C_i^*)_{i \in \mathcal{I}}$ with $\sum_i C_i^* = E$ iff $\langle \psi^*, E \rangle \neq 0$ and

$$V_i(C_i) \geq V_i(C_i^*), C_i \in \tilde{\mathcal{X}}_+ \text{ for all } i \in \mathcal{I} \text{ implies } \left\langle \psi^*, \sum_i C_i \right\rangle \geq \langle \psi^*, E \rangle .$$

We note the following

Proposition 2.5 *If an allocation $(C_i^*)_{i \in \mathcal{I}}$ with $\sum_i C_i^* = E$ is supported by $\psi^* \in L^1(\mathbb{P} \otimes E)$, then $\psi^* > 0$ $\mathbb{P} \otimes dE$ -a.e.*

PROOF : Suppose to the contrary that $\mathbb{P} \otimes dE[\psi^* \leq 0] > 0$. Then, by strict monotonicity and continuity of preferences, we may choose $\epsilon > 0$ small enough such that the allocation defined by

$$dC_i \triangleq (1 - \epsilon) dC_i^* + 1_{\{\psi^* \leq 0\}} dE \quad (i \in \mathcal{I})$$

is preferred to $(C_i^*)_{i \in \mathcal{I}}$. Thus, the supporting property of ψ^* implies

$$\langle \psi^*, E \rangle \leq \left\langle \psi^*, \sum_i C_i \right\rangle = (1 - \epsilon) \langle \psi^*, E \rangle + |\mathcal{I}| \mathbb{E} \int_0^T \psi^* 1_{\{\psi^* \leq 0\}} dE .$$

Hence, we obtain

$$|\mathcal{I}| \mathbb{E} \int_0^T \psi^* 1_{\{\psi^* \leq 0\}} dE \geq \epsilon \langle \psi^*, E \rangle > 0,$$

a contradiction to the obvious relation $\mathbb{E} \int_0^T \psi^* 1_{\{\psi^* \leq 0\}} dE \leq 0$. \square

Now, we may prove

Lemma 2.6 *The Lagrange multiplier ψ_λ supports its associated λ -efficient allocation $(C_i^\lambda)_{i \in \mathcal{I}}$. Moreover, any other optional price density $\psi \in L^1(\mathbb{P} \otimes dE)$ with this property is of the form*

$$\psi = \max_i \{k_i \nabla V_i(C_i^\lambda)\} \quad \mathbb{P} \otimes dE\text{-a.s.}$$

for some constants $k_i \geq 0$ ($i \in \mathcal{I}$), and the λ -efficient allocation satisfies the flat-off conditions

$$\langle \psi - k_i \nabla V_i(C_i^\lambda), C_i^\lambda \rangle = 0 \quad (i \in \mathcal{I}).$$

PROOF : In order to show that ψ_λ supports $(C_i^\lambda)_{i \in \mathcal{I}}$, take an allocation $(C_i)_{i \in \mathcal{I}} \in \tilde{\mathcal{X}}_+^{\mathcal{I}}$ with $V_i(C_i) \geq V_i(C_i^\lambda)$. Concavity of V_i yields

$$0 \leq \sum_i \lambda_i \{V_i(C_i) - V_i(C_i^\lambda)\} \leq \sum_i \lambda_i \langle \nabla V_i(C_i^\lambda), C_i - C_i^\lambda \rangle .$$

By properties (ii) and (iii) of the efficient allocation $(C_i^\lambda)_{i \in \mathcal{I}}$ (Lemma 2.1), the latter quantity is

$$\leq \sum_i \langle \psi_\lambda, C_i - C_i^\lambda \rangle = \left\langle \psi_\lambda, \sum_i C_i - E \right\rangle.$$

Hence, $\langle \psi_\lambda, \sum_i C_i \rangle \geq \langle \psi_\lambda, E \rangle$ which is the claimed supporting property.

For the second part of the lemma, suppose that $\psi \in L^1(\mathbb{P} \otimes dE)$ is optional and supports the allocation $(C_i^\lambda)_{i \in \mathcal{I}}$. We only need to show that there are nonnegative constants k_i ($i \in \mathcal{I}$) such that

$$(1) \quad k_i \nabla V_i(C_i^\lambda) \leq \psi \quad \mathbb{P} \otimes dE\text{-a.e. and } \langle \psi - k_i \nabla V_i(C_i^\lambda), C_i^\lambda \rangle = 0$$

for every $i \in \mathcal{I}$. To this end, put

$$k_i \triangleq \mathbb{P} \otimes dE\text{-ess inf } \frac{\psi}{\nabla V_i(C_i^\lambda)} \quad (i \in \mathcal{I}).$$

Obviously, $k_i \geq 0$ for every $i \in \mathcal{I}$ and, of course, the first condition in (1) is satisfied. To verify the second condition for $i \in \mathcal{I}$, let us distinguish the cases $\langle \psi, C_i^\lambda \rangle = 0$ and $\langle \psi, C_i^\lambda \rangle > 0$.

In the first case, we may conclude from Proposition 2.5 that $C_i^\lambda = 0$. Thus, the second condition in (1) is satisfied trivially. For the case $\langle \psi, C_i^\lambda \rangle > 0$, we prove below that C_i^λ maximizes utility over all consumption plans $C_i \in \tilde{\mathcal{X}}_+$ with $\langle \psi, C_i \rangle \leq \langle \psi, C_i^\lambda \rangle$. By arguments similar to the proof of Theorem 3.3 in Bank and Riedel (1999), one then deduces the validity of the second condition in (1) also in this case.

To obtain the claimed optimality of C_i^λ , it suffices to show that any $C_i \in \tilde{\mathcal{X}}_+$ with $V_i(C_i) > V_i(C_i^\lambda)$ must satisfy $\langle \psi, C_i \rangle > \langle \psi, C_i^\lambda \rangle$. Note first that, for such a C_i , we also have $V_i((1 - \epsilon)C_i) > V_i(C_i^\lambda)$ for any sufficiently small $\epsilon > 0$ by continuity V_i . Consider the allocation $(\tilde{C}_j)_{j \in \mathcal{I}}$ defined by

$$\tilde{C}_j \triangleq \begin{cases} (1 - \epsilon)C_i & \text{for } j = i, \\ C_j^\lambda & \text{otherwise.} \end{cases}$$

This allocation is preferred to $(C_i^\lambda)_{i \in \mathcal{I}}$ and, thus, the supporting property of ψ yields

$$\langle \psi, E \rangle \leq \left\langle \psi, \sum_j \tilde{C}_j \right\rangle = \langle \psi, E \rangle + \langle \psi, (1 - \epsilon)C_i - C_i^\lambda \rangle.$$

This implies

$$0 < \langle \psi, C_i^\lambda \rangle \leq \langle \psi, (1 - \epsilon)C_i \rangle < \langle \psi, C_i \rangle$$

as we wanted to show. □

3 Existence of Equilibria

After these technical preliminaries, we are now in a position to prove existence of equilibria for intertemporal consumption.

Proof of Theorem 1.5 We start by defining the correspondence \mathcal{G} to which Kakutani's fixed point theorem will be applied. To this end, let, for any $\lambda \in \Lambda$, $\mathcal{S}(\lambda)$ denote the set of all optional price densities $\psi \in L^1(\mathbb{P} \otimes dE)$ which support the allocation C^λ and which, in addition, satisfy

$$(2) \quad \frac{b}{|\mathcal{I}| \langle B, E \rangle} \leq \psi \leq \frac{B}{\langle B, E \rangle} \quad \mathbb{P} \otimes dE\text{-a.e.} \quad \text{and} \quad \langle \psi, C_i^\lambda \rangle \leq \lambda_i \quad \text{for every } i \in \mathcal{I}.$$

Here, b and B are the optional gradient bounds introduced in Assumption 2.

We now define the correspondence

$$(3) \quad \mathcal{G}(\lambda) \triangleq \left\{ (\lambda_i + \langle \psi, E_i - C_i^\lambda \rangle)_{i \in \mathcal{I}} \mid \psi \in \mathcal{S}(\lambda) \right\} \quad (\lambda \in \Lambda).$$

In Proposition 3.1 below, we show that indeed \mathcal{G} satisfies the conditions required for Kakutani's theorem. Hence, \mathcal{G} has a fixed point $\lambda^* \in \Lambda$. Let $C^* \triangleq C^{\lambda^*}$ and note that, by definition of \mathcal{G} , there is $\psi^* \in \mathcal{S}(\lambda^*)$ such that $\langle \psi^*, E_i \rangle = \langle \psi^*, C_i^* \rangle$ for every $i \in \mathcal{I}$. As ψ^* supports the efficient allocation C^* , this gives us existence of a quasi-equilibrium in the auxiliary economy where consumption spaces are given by the order ideal $\tilde{\mathcal{X}}_+ \subset \mathcal{X}_+$.

Now, recall from Lemma 2.6 that, as the density ψ^* induces a price functional supporting the allocation $(C_i^*)_{i \in \mathcal{I}}$, it must take the form

$$(4) \quad \psi^* = \max_i \{ k_i^* \nabla V_i(C_i^*) \}$$

for some constants $k_i^* \geq 0$. Note that, via the right side of (4), ψ^* allows a canonical extension to a strictly positive, bounded and optional process on the whole time interval $[0, T]$. This process induces, thus, a price functional $\langle \psi^*, \cdot \rangle$ on the 'large' consumption space \mathcal{X}_+ .

Let us next show that, in conjunction with $(C_i^*)_{i \in \mathcal{I}}$, this functional $\langle \psi^*, \cdot \rangle$ defines a true Arrow–Debreu–equilibrium for the 'large' economy where consumption spaces are given by \mathcal{X}_+ . To this end, fix $i \in \mathcal{I}$ and consider a plan $C_i \in \mathcal{X}_+$ which is strictly preferred to C_i^* , i.e., assume $V_i(C_i) > V_i(C_i^*)$. We have to show that $\langle \psi^*, C_i \rangle > \langle \psi^*, C_i^* \rangle = \langle \psi^*, E_i \rangle$. In fact, we have

$$0 > V_i(C_i^*) - V_i(C_i) \geq \langle \nabla V_i(C_i^*), C_i^* - C_i \rangle \geq \langle \psi^*, C_i^* - C_i \rangle / k_i^*,$$

where the last estimate is due to $k_i^* \nabla V_i(C_i^*) \leq \psi^*$ and to the flat-off condition in Lemma 2.6. Hence, any plan which is strictly preferred to C_i^* violates the investor's budget constraint. \square

It remains to establish

Proposition 3.1 *The correspondence \mathcal{G} defined by (3) satisfies the conditions of Kakutani's fixed point theorem:*

- (i) *For every $\lambda \in \Lambda$, $\mathcal{G}(\lambda)$ is a non-empty convex subset of Λ .*
- (ii) *\mathcal{G} is upper hemi-continuous, i.e.,*

$$\{(\lambda, g) \mid \lambda \in \Lambda, g \in \mathcal{G}(\lambda)\}$$

is closed in $\Lambda \times \Lambda$.

PROOF : We adopt the notation from the preceding proof.

1. Let us focus on assertion (i) and note first that $\mathcal{G}(\lambda)$ is nonempty for every $\lambda \in \Lambda$. Indeed, we know from Lemma 2.6 that ψ_λ supports the allocation C^λ . Clearly, this property is inherited by every positive multiple of ψ_λ . Moreover, we have

$$\frac{b}{|\mathcal{I}|} \leq \psi_\lambda \leq B$$

such that $\psi \triangleq \psi_\lambda / \langle B, E \rangle$ obviously satisfies the first constraint in (2). It also satisfies the second constraint, since by Lemma 2.1 (iii)

$$\langle \psi_\lambda, C_i^\lambda \rangle = \lambda_i \langle \nabla V_i(C_i^\lambda), C_i^\lambda \rangle \leq \lambda_i \langle B, E \rangle .$$

This shows that $\mathcal{S}(\lambda)$ and, hence, also $\mathcal{G}(\lambda)$ is nonempty.

Convexity of $\mathcal{G}(\lambda)$ follows from convexity of $\mathcal{S}(\lambda)$. Moreover, any $g \in \mathcal{G}(\lambda)$ satisfies $g_i \geq 0$ for every $i \in \mathcal{I}$ because of the second constraint in (2). In addition, $\sum_i g_i = 1$ by Lemma 2.1 (i). Hence, we have in fact $\mathcal{G}(\lambda) \subset \Lambda$ which completes the proof of assertion (i).

2. To prove assertion (ii), let $\lambda^n \in \Lambda$ and $g^n \in \mathcal{G}(\lambda^n)$ ($n = 1, 2, \dots$) converge to λ^0 and g^0 , respectively. We have to show that $g^0 \in \mathcal{G}(\lambda^0)$. Put $C^n \triangleq C^{\lambda^n}$ and let $\psi_n \in \mathcal{S}(\lambda^n)$ be such that

$$(5) \quad g_i^n = \lambda_i^n + \langle \psi_n, E_i - C_i^n \rangle \quad (i \in \mathcal{I}).$$

Due to condition (2), the sequence $(\psi_n, n = 1, 2, \dots)$ is dominated by the $\mathbb{P} \otimes dE$ -integrable process $B / \langle B, E \rangle$. In particular, it is uniformly integrable and, by the Dunford–Pettis–theorem, there is a subsequence, again denoted by (ψ_n) , which converges weakly to some ψ in $L^1(\mathbb{P} \otimes dE)$.

We shall show that ψ belongs to $\mathcal{S}(\lambda^0)$ and satisfies

$$(6) \quad g_i^0 = \lambda_i^0 + \langle \psi, E_i - C_i^0 \rangle \quad \text{for every } i \in \mathcal{I}.$$

Of course, this will yield assertion (ii). Our argument is based on the following result

$$(7) \quad \lim_n \langle \psi_n, C_i^n \rangle = \langle \psi, C_i^0 \rangle \quad \text{for every } i \in \mathcal{I}$$

which goes back to Bewley (1969) and which will be established in Step 3 of this proof.

As a first application of Bewley's claim, we note that (6) holds true. Indeed, granted (7), this follows immediately by letting $n \uparrow \infty$ in (5).

Similarly, we show that ψ satisfies the second condition in (2) for $\lambda = \lambda^0$. Indeed as $\psi_n \in \mathcal{S}(\lambda^n)$ by definition, we know that each ψ_n satisfies $\langle \psi_n, C_i^n \rangle \leq \lambda_i^n$ for every $i \in \mathcal{I}$. Given relation (7), we may pass to the limit $n \uparrow \infty$ to obtain the desired inequalities

$$\langle \psi, C_i^0 \rangle \leq \lambda_i^0 \quad (i \in \mathcal{I}).$$

The first condition in (2) is stable with respect to weak convergence in $L^1(\mathbb{P} \otimes dE)$ and is thus inherited by ψ from ψ_n ($n = 1, 2, \dots$).

Concerning the supporting property of ψ , note first that, from $\psi \geq b/(|\mathcal{I}| \langle B, E \rangle)$ it follows that

$$\langle \psi, E \rangle \geq \frac{\langle b, E \rangle}{|\mathcal{I}| \langle B, E \rangle} > 0.$$

Hence, in order to verify $\psi \in S(\lambda^0)$, it only remains to show that, under the price density ψ , every allocation $C \in \tilde{\mathcal{X}}_+^{\mathcal{I}}$ which is preferred to C^0 must have a higher aggregate price than C^0 . By monotonicity of preferences, it suffices to consider a *strictly* preferred allocation C in the sense that $V_i(C_i) > V_i(C_i^0)$ for every $i \in \mathcal{I}$. Due to Corollary 2.3 such an allocation C is also strictly preferred to C^n when n is large enough. As $\psi_n \in \mathcal{S}(\lambda^n)$ by assumption, each C^n is supported by ψ_n ($n = 1, 2, \dots$). We thus obtain that, for n sufficiently large,

$$\left\langle \psi_n, \sum_i C_i \right\rangle \geq \langle \psi_n, E \rangle.$$

Since (ψ_n) converges weakly in $L^1(\mathbb{P} \otimes dE)$ to ψ and $\sum_i C_i$ and E belong to the order ideal $\tilde{\mathcal{X}}_+$, it follows that

$$\left\langle \psi, \sum_i C_i \right\rangle \geq \langle \psi, E \rangle.$$

This shows that ψ indeed supports C^0 and, therefore, completes the proof of assertion (ii).

3. We still have to prove Bewley's claim (7). We follow his argument and note first that the claim already follows from the seemingly weaker assertion

$$(8) \quad \limsup_n \langle \psi_n, C_i^n \rangle \leq \langle \psi, C_i^0 \rangle \quad \text{for every } i \in \mathcal{I}.$$

Indeed, the aggregation property of allocations and claim (8) imply

$$(9) \quad \langle \psi, E \rangle \geq \limsup_n \sum_i \langle \psi_n, C_i^n \rangle = \limsup_n \langle \psi_n, E \rangle .$$

Due to the weak convergence $\psi_n \rightarrow \psi$, the last term is again equal to $\langle \psi, E \rangle$. Hence, we must have equality everywhere in (9) and claim (7) follows.

In order to establish (8), fix $i \in \mathcal{I}$ and set $C_i^\epsilon \triangleq C_i^0 + \epsilon E$ for $\epsilon > 0$. By monotonicity of preferences, we have $V_i(C_i^\epsilon) > V_i(C_i^0)$. Due to Corollary 2.3, we also have

$$V_i(C_i^\epsilon) > V_i(C_i^m) \quad \text{for large } n.$$

Since ψ_n supports the allocation C^n , it follows that for such n

$$\left\langle \psi_n, C_i^\epsilon + \sum_{j \neq i} C_j^m \right\rangle \geq \langle \psi_n, E \rangle$$

or, equivalently,

$$\langle \psi_n, C_i^\epsilon \rangle \geq \langle \psi_n, C_i^m \rangle .$$

Let n tend to infinity and use the weak convergence $\psi_n \rightarrow \psi$ to conclude

$$\langle \psi, C_i^\epsilon \rangle \geq \limsup_n \langle \psi_n, C_i^m \rangle .$$

Now, claim (8) follows from letting $\epsilon \rightarrow 0$ in the preceding estimate. □

4 Structure of Equilibrium Prices

Having established existence of an equilibrium $((C_i^*)_{i \in \mathcal{I}}, \psi^*)$, it is natural to ask, whether the induced equilibrium price functional $\langle \psi^*, \cdot \rangle$ is continuous on our consumption space $(\mathcal{X}_+, d_{\mathcal{X}_+})$. Indeed, this is desirable from an economic point of view, since ‘similar’ consumption plans in \mathcal{X}_+ should have ‘similar’ prices. We show in this section that under two additional assumptions, we indeed have this kind of continuity of the price functional.

Our first condition is

Assumption 3 *The filtration $\mathbb{F} = (\mathcal{F}_t, 0 \leq t \leq T)$ is quasi left-continuous.*

This is an assumption on the way new information is revealed to the agents. It is satisfied if, e.g., the filtration \mathbb{F} is generated by a Brownian motion or a Poisson process. Economically, an information flow corresponds to a quasi left-continuous filtration if ‘information surprises’ (in the sense of Hindy and Huang (1992)) occur only at times which cannot be predicted. An earthquake in New York (rather than San Francisco) is an example. The announcement of a policy change of the Federal reserve is an example for an information surprise which occurs at a time known in advance.

The second assumption allows us to use stochastic calculus:

Assumption 4 For every $C \in \mathcal{X}_+$, each utility gradient $\nabla V_i(C)$ ($i \in \mathcal{I}$) is a bounded (\mathbb{P}, \mathbb{F}) -semimartingale with a continuous compensator of bounded variation.

Clearly, this assumption is satisfied, e.g., if all agents have Hindy–Huang–Kreps preferences as described in Section 1.3.

Theorem 4.1 Under Assumptions 3 and 4, every equilibrium price functional

$$C \mapsto \langle \psi^*, C \rangle \quad (C \in \mathcal{X}_+)$$

is continuous on $(\mathcal{X}_+, d_{\mathcal{X}_+})$.

PROOF :

1. As an equilibrium price process, ψ^* clearly supports its associated equilibrium allocation $C^* \triangleq (C_i^*)_{i \in \mathcal{I}}$. By Lemma 2.1, it thus takes the form

$$\psi^* = \max_i \{k_i^* \nabla V_i(C_i^*)\}$$

for suitable constants $k_i^* \geq 0$ ($i \in \mathcal{I}$).

Due to Assumption 4, each process $\phi_i \triangleq k_i^* \nabla V_i(C_i^*)$ ($i \in \mathcal{I}$) allows a Doob–Meyer decomposition $\phi_i = M_i + A_i$ into a local martingale M_i and a continuous compensator A_i of bounded variation.

Moreover, defined as the pointwise maximum of the bounded semimartingales ϕ_i , the process ψ is a bounded semimartingale, too. Hence, it can be decomposed in the form $\psi = M + A$ where M is a local martingale and A is a predictable RCLL-process of bounded variation.

In particular, there is a localizing sequence of stopping times T_m ($m = 1, 2, \dots$) such that each of the stopped processes $M^{T_m}, M_i^{T_m}$ ($i \in \mathcal{I}$) is a (uniformly integrable) martingale on $[0, T]$ and such that eventually $T_m(\omega) = T$ for \mathbb{P} -a.e. $\omega \in \Omega$.

2. We claim that A almost surely has continuous paths. In order to prove this, it suffices to show that almost surely $A(S) = A(S-)$ for every *predictable* stopping time $S \leq T$ because both processes $(A(t), (0 \leq t \leq T))$ and $(A(t-), (0 \leq t \leq T))$ are predictable; cf., e.g., Rogers and Williams (1987), Lemma VI.19.2.

Now, recall that, granted quasi-left continuity of the underlying filtration, every uniformly integrable martingale almost surely does not jump at predictable times; see, e.g., Theorem VI.18.1 in Rogers and Williams (1987).

We apply this observation first to the martingales $M_i^{T_m}$ and obtain that $\Delta M_i(S) = 0$ on $\{S \leq T_m\}$ for every $i \in \mathcal{I}$. Since, in addition, each A_i is continuous, this yields $\Delta \phi_i(S) = 0$ on $\{S \leq T_m\}$ for every $i \in \mathcal{I}$. For $m \uparrow +\infty$, this entails $\Delta \phi_i(S) = 0$ ($i \in \mathcal{I}$) and, consequently, also $\Delta \psi(S) = 0$ almost surely.

Applying the above observation to the stopped process M^{T_m} shows that also $\Delta M(S) = 0$ on $\{S \leq T_m\}$. Letting $m \uparrow \infty$ we obtain $\Delta M(S) = 0$ almost surely. Together with $\Delta \psi(S) = 0$, this implies $\Delta A(S) = 0$ and we are done.

3. We are now in a position to prove the asserted continuity of the price functional $C \mapsto \langle \psi^*, C \rangle$. To this end, let C^n ($n = 1, 2, \dots$) converge to C^0 in the metric $d_{\mathcal{X}_+}$. Thus, we have $L^1(\mathbb{P})$ -convergence of $C^n(T)$ to $C^0(T)$ and weak*-convergence in probability of the measures dC^n to dC^0 . By the usual subsequence argument, we may assume without loss of generality that both convergences hold true even almost surely.

Note that the local martingale M is locally bounded since it is the difference of the bounded process ψ and the continuous process A . Thus, we may assume that our localizing sequence (T_m) is such that each M^{T_m} ($m = 1, 2, \dots$) is a bounded martingale.

For ease of notation, we put $\psi \triangleq \psi^*$. For every $m = 1, 2, \dots$ we have

$$\begin{aligned} |\langle \psi, C^n \rangle - \langle \psi, C^0 \rangle| &\leq \left| \mathbb{E} \int_0^T [\psi - \psi^{T_m}] (dC^n - dC^0) \right| + \left| \mathbb{E} \int_0^T \psi^{T_m} (dC^n - dC^0) \right| \\ &\leq \mathbb{E} \int_{T_m}^T |\psi - \psi^{T_m}| (dC^0 + dC^n) \\ &\quad + \left| \mathbb{E} \int_0^T M^{T_m} (dC^n - dC^0) \right| + \left| \mathbb{E} \int_0^T A^{T_m} (dC^n - dC^0) \right|. \end{aligned}$$

Let us denote the preceding three summands by I , II , and III , respectively. For the first summand, we have

$$I \leq \|\psi\|_\infty \mathbb{E} [(C^n(T) + C^0(T)) 1_{\{T_m < T\}}].$$

As $C^n(T) \rightarrow C^0(T)$ in $L^1(\mathbb{P})$ by assumption, dominated convergence yields

$$\limsup_n I \leq 2\|\psi\|_\infty \mathbb{E} [C^0(T) 1_{\{T_m < T\}}].$$

Using the martingale property of M^{T_m} , we may rewrite the second summand in the form

$$II = |\mathbb{E} [M(T_m)(C^n(T) - C^0(T))]|.$$

Thus,

$$\limsup_n II \leq \|M(T_m)\|_\infty \limsup_n \mathbb{E} |C^n(T) - C^0(T)| = 0.$$

Finally, note that, due to the continuity of A , we have

$$(10) \quad \int_0^T A^{T_m} dC^n \rightarrow \int_0^T A^{T_m} dC^0 \quad (n \uparrow \infty)$$

almost surely. Moreover, we have

$$\left| \int_0^T A^{T_m} dC^n \right| \leq \left\| \sup_{0 \leq t \leq T} |A^{T_m}(t)| \right\|_\infty C^n(T).$$

Once again we use the $L^1(\mathbb{P})$ -convergence $C^n(T) \rightarrow C^0(T)$ and deduce that the right side of this estimate defines a uniformly integrable family of random variables parametrized by n . Hence, by Lebesgue's Theorem, we obtain that the convergence (10) holds true also in $L^1(\mathbb{P})$. For the third summand, this gives us

$$\limsup_n III = 0.$$

Summing up, we find that, for every $m = 1, 2, \dots$,

$$(11) \quad \begin{aligned} \limsup_n |\langle \psi, C^n \rangle - \langle \psi, C^0 \rangle| &\leq \limsup_n I + \limsup_n II + \limsup_n III \\ &\leq 2\|\psi\|_\infty \mathbb{E} [C^0(T) 1_{\{T_m < T\}}] + 0 + 0. \end{aligned}$$

Letting $m \uparrow \infty$ in (11), we get by dominated convergence that indeed

$$\limsup_n |\langle \psi, C^n \rangle - \langle \psi, C^0 \rangle| \leq 2\|\psi\|_\infty \mathbb{E} [C^0(T) 1_{\cap_m \{T_m < T\}}] = 0$$

because $T_m(\omega) = T$ eventually for \mathbb{P} -a.e. $\omega \in \Omega$.

□

Properties of Equilibrium Price Processes We study now the properties of equilibrium price processes in more detail.

Being the weighted maximum of utility gradients, the equilibrium price process is a semimartingale if gradients are semimartingales. This is an important property because it provides an equilibrium foundation for the application of stochastic calculus in mathematical finance.

Going a step further, we see from the preceding proof that the bounded variation part of the equilibrium price process is continuous if gradients' bounded variation parts are continuous.

A fundamental question is whether this bounded variation part is even absolutely continuous because then an interest rate exists. The gradients of Hindy–Huang–Kreps utility functionals have such a nice representation. Hence, in a one consumer world, the Hindy–Huang–Kreps approach guarantees the existence of an interest rate. Hindy and Huang (1992) even suggest that also with heterogeneous agents, the equilibrium price process would have this nice property. However, this need not always be the case. From the Tanaka formula, it follows that the maximum of semimartingales whose finite variation part is absolutely continuous can be decomposed into a local martingale, an absolutely continuous part of bounded variation and a part which depends on local time of the gradients. In general, local times are not absolutely continuous. Hence, the equilibrium price process possibly does not belong to the dual suggested by Hindy and Huang (1992). Moreover, our characterization of supporting price functionals (Lemma 2.6) shows that, in general, there may be no equilibrium whose price process is contained in the Hindy–Huang dual.

Local time arises in the decomposition of the equilibrium price process whenever the identity of the agent whose gradient determines the price changes. Hence, whenever one agent stops consuming and another restarts, local time may appear. Such a phenomenon has already been remarked in the time-additive setting by Karatzas, Lehoczky, and Shreve (1991). For finance theory, this has the implication that the money market account contains a singular component. The detailed consequences for finance theory remain to be studied in future work.

References

- BANK, P., AND F. RIEDEL (1999): “Optimal Consumption Choice Under Uncertainty with Intertemporal Substitution,” Discussion paper 71, Sonderforschungsbereich 373, Humboldt University Berlin.
- (2000): “Optimal Consumption Choice with Intertemporal Substitution,” mimeo, Humboldt University Berlin.
- BEWLEY, T. (1969): “A Theorem on the Existence of Competitive Equilibria in a Market with a Finite Number of Agents and Whose Commodity Space is L^∞ ,” in *Studies in Economic Theory*, ed. by M. Khan, and N. Yannelis, vol. 1, pp. 74–101. Springer, 1991, New York, Heidelberg.
- BREEDEN, D. (1979): “An Intertemporal Asset Pricing Model with Stochastic Consumption and Investment Opportunities,” *Journal of Financial Economics*, 7, 265–296.
- DUFFIE, D., AND L. EPSTEIN (1992): “Stochastic Differential Utility,” *Econometrica*, 60, 353–394.
- DUFFIE, D., AND C. SKIADAS (1994): “Continuous-Time Security Pricing, a Utility Gradient Approach,” *Journal of Mathematical Economics*, 23, 107–131.
- DUFFIE, D., AND W. ZAME (1989): “The Consumption-Based Capital Asset Pricing Model,” *Econometrica*, 57, 1274–1298.
- HINDY, A., AND C.-F. HUANG (1992): “Intertemporal Preferences for Uncertain Consumption: a Continuous-Time Approach,” *Econometrica*, 60, 781–801.
- HINDY, A., C.-F. HUANG, AND D. KREPS (1992): “On Intertemporal Preferences in Continuous Time — the Case of Certainty,” *Journal of Mathematical Economics*, 21, 401–440.
- HINDY, A., C.-F. HUANG, AND S.-H. ZHU (1997): “Optimal Consumption and Portfolio Rules with Durability and Habit Formation,” *Journal of Economic Dynamics and Control*, 21, 525–550.

- KABANOV, Y. (1999): “Hedging and Liquidation under Transaction Costs in Currency Markets,” *Finance and Stochastics*, 3, 237–248.
- KARATZAS, I., J. P. LEHOCZKY, AND S. E. SHREVE (1991): “Equilibrium Models with Singular Asset Prices,” *Mathematical Finance*, 1(3), 11–29.
- KOMLÓS, J. (1967): “A Generalization of a Problem of Steinhaus,” *Acta Math. Acad. Sci. Hung.*, 18, 217–229.
- MAS-COLELL, A., AND S. RICHARD (1991): “A New Approach to the Existence of Equilibria in Vector Lattices,” *Journal of Economic Theory*, 53, 1–11.
- MAS-COLELL, A., AND W. ZAME (1991): “Equilibrium Theory in Infinite Dimensional Spaces,” in *Handbook of Mathematical Economics*, ed. by W. Hildenbrand, and H. Sonnenschein, vol. 4, pp. 1835–1898. Elsevier Science Publishers B.V., North-Holland, Amsterdam.
- MERTON, R. C. (1971): “Optimum Consumption and Portfolio Rules in a Continuous-Time Model,” *Journal of Economic Theory*, 3, 373–413.
- ROGERS, L., AND D. WILLIAMS (1987): *Diffusions, Markov Processes, and Martingales*, vol. 2 of *Series in Probability and Mathematical Statistics*. Wiley, Chichester.

PETER BANK

Institut für Mathematik
Humboldt-Universität zu Berlin
Unter den Linden 6
D-10099 Berlin, Germany
E-mail: pbank@mathematik.hu-berlin.de

FRANK RIEDEL

Institut für Wirtschaftstheorie I
Humboldt-Universität zu Berlin
Spandauer Straße 1
D-10178 Berlin, Germany
E-mail: riedel@odin.wiwi.hu-berlin.de