
**12. Practice sheet for the lecture:
Combinatorics (DS I)**

Felsner/Heldt, Knauer
28. June

Delivery date: 4. –8. July

<http://www.math.tu-berlin.de/~felsner/Lehre/dsI11.html>

- (1) For which of the parameter sets does a design exist? Either show that there is no design or present one.
 - (a) $t = 4, k = 7, v = 13$ and $\lambda = 2$.
 - (b) $t = 2, k = 7, v = 36$ and $\lambda = 1$.
 - (c) $t = 2, k = 5, v = 125$ and $\lambda = 1$.
 - (d) $t = 1, k = 4, v = 124$ and $\lambda = 1$.
- (2) Let (V, \mathcal{B}) be a $S_\lambda(t, k, v)$ design. Let $p \in V$ and $\mathcal{B}^p := \{B : p \notin B \in \mathcal{B}\}$ be the set of blocks, which do not contain p . Show that $(V \setminus \{p\}, \mathcal{B}^p)$ is a design. What are its parameters?
- (3) Let $(V, \mathcal{B}) = S(2, n + 1, n^2 + n + 1)$ be a projective plane and fix $B \in \mathcal{B}$. Show that $(V \setminus B, \{C \setminus B \mid C \in (\mathcal{B} \setminus \{B\})\})$ is a $S(2, n, n^2)$ design.
- (4) Prove Fisher's proposition, which states that every $S_\lambda(t, k, v)$ design (V, \mathcal{B}) with $t \geq 2$ fulfills $|V| \leq |\mathcal{B}|$ (Hint: Use the adjacency matrix $A \in \mathbb{R}^{|V| \times |\mathcal{B}|}$ with $a_{v,B} = 1$ if $v \in B$ and $a_{v,B} = 0$ otherwise and consider the rank of $A \cdot A^T$).
- (5) Let (V, \mathcal{B}) be a design, $I, J \subseteq V$ with $I \cap J = \emptyset$ and $|I| = i, |J| = j$ such that $i + j \leq t$ and $k < v$. Let $\lambda_{I,J} = \#\{B \in \mathcal{B} \mid I \subseteq B \text{ and } J \cap B = \emptyset\}$.
 - (a) Show that $\lambda_{I,J}$ does only depend on i and j and not on I and J , i.e. $\lambda_{i,j} := \lambda_{I,J}$ is well defined.
 - (b) Compute all $\lambda_{i,j}$ for the $S_6(3, 5, 10)$ design from the lecture.
 - (c) Prove $\lambda_{i,j} = \lambda_{i+1,j} + \lambda_{i,j+1}$ for $i + j < t$.
 - (d) Prove $\lambda_{i,j} = \sum_{r=0}^j (-1)^r \binom{j}{r} \lambda_{i+r,0}$.