
1. Practice sheet for the lecture:
Combinatorics (DS I)

Felsner/ Kleist
18. April 2017

Due dates: 25.,27. April

<http://www.math.tu-berlin.de/~felsner/Lehre/dsI17.html>

- (1) A spider has a sock and a shoe for each of his eight feet. In how many different ways can he put on his shoes and socks, assuming that on each foot he has to put on the sock first?
- (2) The squares of a 4×7 chessboard are coloured arbitrarily black and white (i.e. there are $2^{4 \cdot 7}$ colourings). Show that there is a $i \times j$ -rectangle with $i \geq 2$ and $j \geq 2$, such that all four of its corners have the same colour. Is the same true for a 4×6 chessboard?
- (3) Consider a chess tournament of n players, each playing against every other participant. Show that at each point of time during the tournament there exist at least two players, having finished the same number of games.
- (4) A sequence of numbers a_1, a_2, \dots, a_n is called *unimodal*, if there exists an $m \in [n]$, such that $a_i \leq a_{i+1}$ for all $i < m$ and $a_i \geq a_{i+1}$ for all $i \geq m$. Give three different proofs of the unimodality of the sequence $\binom{n}{1}, \binom{n}{2}, \binom{n}{3}, \dots, \binom{n}{n}$ for all $n \in \mathbb{N}$, based on the three given hints:

- (a) Use the definition

$$\binom{n}{k} := \frac{n!}{k! \cdot (n-k)!}.$$

- (b) Consider the recursive definition

$$\binom{n}{k} := \binom{n-1}{k-1} + \binom{n-1}{k} \text{ and } \binom{n}{0} = \binom{n}{n} = 1,$$

based on Pascale's triangle.

- (c) Use the bijection between $\binom{n}{k}$ and the number of subsets of $[n]$, having k elements.
- (5) Permutations

- (a) Give an alternative proof for the generating function of derangements (6)

$$D(z) := \sum \frac{d(n)}{n!} z^n = \frac{e^{-z}}{1-z}$$

by manipulating $D(z)e^z$ and using identity (2a) from the lecture.

- (b) Prove of the following identity:

$$d(n) = n \cdot d(n-1) + (-1)^n$$

- (c) How many permutations in S_n are derangements and involutions?
(Let X be a set. A function $f : X \rightarrow X$ is an *involution* if $f(f(x)) = x \forall x \in X$.)
- (d) What is the expected number of fixed points of a uniform random permutation?