

# Differentialgeometrie II

## Übungsblatt 9

Due on Wednesday 28.01.2009

### 1 Aufgabe

Let  $\phi \in \Lambda^r(\mathbb{R}^n)$ . Show that  $\phi$  is closed iff  $\int_{\partial B} \phi = 0$  for any  $r + 1$  dimensional submanifold  $B$  of  $\mathbb{R}^n$ .

### 2 Aufgabe

Let  $M$  be an  $n$ -dimensional oriented manifold, with orientation determined by  $\Omega \in \Lambda^n(M)$ .

- Show that if  $M$  is Riemannian, the metric determines uniquely a vector field  $X$  defined at each point  $p \in \partial M$  and such that  $X_p$  is orthogonal to  $T_p \partial M$ , inward pointing (cfr. remark VI.4.5) and of unit length.
- Show that in the general case where  $M$  is not Riemannian, it is possible to find a vector field  $X$  defined at each point  $p \in \partial M$  such that it is inward pointing (use partition of unity).
- Show that  $(-1)^{n-1} \iota_X(\Omega) \in \Lambda^{n-1}(\partial M)$  determines on  $\partial M$  the same orientation as in theorem VI.4.4.

### 3 Aufgabe

Let  $S^n$  be the unit sphere, viewed as a submanifold of  $\mathbb{R}^{n+1}$ . We endow  $\mathbb{R}^{n+1}$  with standard coordinates  $x^1, \dots, x^{n+1}$  and standard volume form  $\Omega = dx^1 \wedge \dots \wedge dx^{n+1}$ .

Consider  $B_{n+1}$  (the unit ball) and define on  $S^n = \partial B_{n+1}$  an orthonormal inward pointing vector field  $X$  as in the first part of problem 2.

Write the expression for  $\omega := (-1)^n \iota_X(\Omega) \in \Lambda^n(S^n)$ .

Consider the map  $\sigma: S^n \rightarrow S^n$  sending  $(x^1, \dots, x^n, x^{n+1})$  to  $(-x^1, \dots, -x^n, -x^{n+1})$ .

Show that  $\sigma^*(\omega) = (-1)^{n+1} \omega$ .

### 4 Aufgabe

Let  $S^n$  be as in the preceding problem.

Consider the equivalence relation  $\sim$  on  $S^n$  defined as  $x \sim y$  iff  $x = \sigma(y)$ , for  $x, y \in S^n$ .

Let  $N := S^n / \sim$  be the quotient manifold and  $\pi: S^n \rightarrow N$  the quotient map. Show that

- $\forall f \in \Lambda^0(N), \sigma^* \circ \pi^*(f) = \pi^*(f) \in \Lambda^0(S^n)$ ;
- $\forall f \in \Lambda^0(N), \sigma^* \circ \pi^*(df) = \pi^*(df) \in \Lambda^1(S^n)$ ;
- $\forall \phi \in \Lambda^r(N), \sigma^* \circ \pi^*(\phi) = \pi^*(\phi) \in \Lambda^r(S^n)$ .

Conclude that for  $n$  even,  $N$  is not orientable.

*Hint:* remember from the preceding problem that  $\sigma^*(\omega) = (-1)^{n+1} \omega$ .